

SCIENCE :

A WEEKLY RECORD OF SCIENTIFIC PROGRESS.

JOHN MICHELS, Editor.

PUBLISHED AT
229 BROADWAY, NEW YORK.
P. O. Box 8838.

SATURDAY, SEPTEMBER 18, 1880.

PROFESSOR ALEXANDER AGASSIZ'S address, delivered in Saunders' Theatre, Cambridge, which we print in full on another page, must be considered one of the most important events of the great Boston meeting.

In his position as Vice-President of the Biological Section nothing could be more appropriate than the expression of his views upon the direction which modern biological research is taking. Animated by his own experience and convictions, his address was a deliberate and able attack upon the prevailing tendency towards too rapid generalization—a tendency which has been increasing during the last fifteen years, and is clearly the outgrowth of the intense desire of modern biologists to break down each and every barrier which obstructs our view in the history of development. Natural and laudable as is the desire to leave no stone unturned in our knowledge of the relationships of the different branches of the animal kingdom, it can only result in the obstruction of future investigators if it is not kept strictly within the limits of the truth. Phylogenetic inquiries add greatly to the zest of study, but should not be carried so far as to hamper or obscure the real end in view, which is, of course, truth and precision of statement, with the line sharply defined between what is actually seen and that which it is inferred ought to be seen.

Prof. Agassiz based his conclusions upon his comprehensive study of the sea-urchin. Stating as a premise the now well-known fact that in their embryological development the modern forms repeat the stages through which their ancestors passed in fossil history, he carefully traced the parallelism in a number of modern and fossil forms, giving an outline of his recent study. The results have been in all cases in positive confirmation of the above premise, and show the very close affiliation of the oldest and most recent forms, in general characters. But while the sea-urchins, with a comparatively

small number of existing species, and with a comparatively complete fossil record, offer a tempting field for speculation, Prof. Agassiz denied his right to group the genera into anything like a complete genealogical tree. "If," he concluded, "when we take one of the most limited groups of the animal kingdom, we find ourselves engaged in a hopeless task, what must be the prospect should we attack the problem of other classes or groups of the animal kingdom, where the species run into thousands, while they number only tens in the case we have attempted to carry out? Shall we say 'ignorbimus' or 'impavidi progrediamur,' and valiantly chase a phantom we can never hope to seize?"

It was hardly to be expected that such an attack as this would pass unnoticed, and in fact, one of the features of the meetings of the Biological Section was a debate growing out of it, which took place on the following day. Prof. Cope had been reading an able paper upon the succession of the extinct Felidæ, pointing to the modifications of the teeth as a basis for forming a complete genetic series. At the close of his paper he called Prof. Agassiz's attention to the fact that here, in the cat family, was an instance leading in quite an opposite direction to that which Prof. Agassiz had assumed in his address the day before. An interesting discussion followed. Prof. Agassiz said he did not object to the grouping of genera into lines of descent where the structural characters were sufficiently homologous, but he did object to regarding such affinity as justifying the introduction of hypothetical links into other parts of the chain, and he did not see that the modifications of a single character, the teeth, warranted the phylogenetic conclusions which Prof. Cope had just reached. Prof. Burt Wilder added that, in his own study upon the pectoral muscles of the dog and other animals, he had found the fallacy of hasty generalization for genetic inferences, drawn from the muscles alone, would widely differ from the facts of actual relationship. Prof. Cope replied that in such questions all must admit that different values should be assigned to different parts of the animal frame, and among the hard parts, of course, he ranked, first, the limbs, then, he said, came the teeth. In justification of his arrangement of the extinct cats into two lines of descent from a common ancestor, he said that the complication of the brains confirmed the history told by the teeth. Prof. Agassiz emphatically repeated his statement of the day before and the discussion closed. We hope this address will be widely circulated and read; if received in the spirit in which Prof. Agassiz intended it, its effect will be admirable. The reaction from the theory of special creation is running strongly in every quarter, and in a day when we find ingenious speculations advanced even in small memoirs, every one must admit the necessity of a more conservative spirit. There is no danger of going into the old and opposite extreme, nor does Prof. Agassiz's address encourage the return movement. It is a re-statement of the old piece of advice—do not attempt to run before you are sure you can walk.

ADDRESS BY ALEXANDER AGASSIZ,

PALEONTOLOGICAL AND EMBRYOLOGICAL DEVELOPMENT.

Since the publication of the "*Poissons Fossiles*" by Agassiz and of the "*Embryologie des Salmonidées*" by Vogt, the similarity, traced by the former between certain stages in the growth of young fishes and the fossil representatives of extinct members of the group, has also been observed in nearly every class of the animal kingdom, and the fact has become a most convenient axiom in the study of paleontological and embryological development. This parallelism, which has been on the one side a strong argument in favor of design in the plan of creation, is now, with slight emendations, doing duty on the other as a newly discovered article of faith in the new biology.

But while in a general way we accept the truth of the proposition that there is a remarkable parallelism between the embryonic development of a group and its paleontological history, yet no one has attempted to demonstrate this, or rather to show how far the parallelism extends. We have up to the present time been satisfied with tracing the general coincidence, or with striking individual cases.

The resemblance between the pupa stage of some Insects and of adult Crustacea, the earlier existence of the latter, and the subsequent appearance of the former in paleontological history, furnished one of the first and most natural illustrations of this parallelism; while theoretically the necessary development of the higher tracheate insects from their early branchiate aquatic ancestors seemed to form an additional link in the chain, and point to the Worms, the representatives of the larval condition of Insects, as a still earlier embryonic stage of the Articulates.

Indeed, there is not a single group of the animal kingdom in which embryology has not played a most important part in demonstrating affinities little suspected before. The development of our frogs, our salamanders, has given us the key to much that was unexplained in the history of Reptiles and Batrachians. The little that has been done in the embryology of Birds has revolutionized our ideas of a class which at the beginning of the century seemed to be the most naturally circumscribed of all. Embryology and paleontology combined have led to the recognition of a natural classification uniting Birds and Reptiles on the one side and Batrachians and Fishes on the other. It is to embryology that we owe the explanation of the affinities of the old Fishes in which Agassiz first recognized the similarity to the embryo of Fishes now living, and by its aid we may hope to understand the relationship of the oldest representatives of the class. It has given us the only explanation of the early appearance of the Cartilaginous Fishes, and of the probable formation of the earliest vertebrate limb from the lateral embryonic fold, still to be traced in the young of the Osseous Fishes of to-day.

Embryology has helped us to understand the changes aquatic animals must gradually undergo in order to become capable of living upon dry land. It has given us pictures of swimming-bladders existing as rudimentary lungs in Fishes with a branchial system; in Batrachians it has shown us the persistence of a branchial system side by side with a veritable lung. We find among the earliest terrestrial Vertebrates, types having manifest affinities with the Fishes on one side and Batrachians on the other, and we call these types Reptiles; but we should nevertheless do so with a reservation, looking to embryology for the true meaning of these half-fledged Reptiles, which lived at the period of transition between an aquatic and terrestrial life, and must therefore always retain an unusual importance in the study of the development of animal life.

When we come to the embryology of the marine Invertebrates, the history of the development of the barnacles is too familiar to be dwelt upon, and I need only allude to the well-known transformations of the Echinoderms, of the Acalephs, Polyps, in fact of every single class of Invertebrates, and perhaps in none more than in the Brachiopods, to show how far-reaching has been the influence of embryology

in guiding us to a correct reading of the relations between the fossils of successive formations. There is scarcely an embryological monograph now published dealing with any of the later stages of growth which does not speak of their resemblance to some type of the group long ago extinct. It has therefore been most natural to combine with the attempts constantly made to establish the genetic sequence between the genera of successive formations, an effort to establish also a correspondence between their paleontological sequence and that of the embryonic stages of development of the same, thus extending the mere similarity first observed between certain stages to a far broader generalization.

It would carry me too far to sketch out, except in a most general way, even for a single class, the agreement known to exist in certain groups between their embryonic development and their paleontological history. It is hinted at in the succession of animal life of any period we may take up, and perhaps cannot be better expressed than by comparing the fauna of any period as a whole with that of following epochs—a zoological system of the Jura, for instance, compared with one made up for the Cretaceous; next, one for the Tertiary, compared with the fauna of the present day. In no case could we find any class of the animal kingdom bearing the same definitions or characterized in the same manner. But apply to this comparison the data obtained from the embryological development of our present fauna, and what a flood of light is thrown upon the meaning of the succession of these apparently disconnected animal kingdoms, belonging to different geological periods, especially in connection with the study of the few ancient types which have survived to the present day from the earliest times in the history of our earth!

Although there is hardly a class of the animal kingdom in which some most interesting parallelism could not be drawn, and while the material for an examination of this parallelism is partially available for the Fishes, Mollusks, Crustacea, Corals, and Crinoids, yet for the illustration and critical examination of this parallelism I have been led to choose to-day a very limited group, that of Sea-urchins, both on account of the nature of the material and of my own familiarity with their development and with the living and extinct species of Echini. The number of living species is not very great—less than three hundred—and the number of fossil species thus far known is not, according to Zittel, more than about two thousand. It is therefore possible for a specialist to know of his own knowledge the greater part of the species of the group. It has been my good fortune to examine all but a few of the species now known to exist, and the collections to which I have had access contain representatives of the majority of the fossil species. Sea-urchins are found in the oldest fossiliferous rocks; they have continued to exist without interruption in all the strata up to the present time. While it is true that our knowledge of the Sea-urchins occurring before the Jurassic period is not very satisfactory, it is yet complete enough for the purposes of the present essay, as it will enable me, starting from the Jurassic period, to call your attention to the paleontological history of the group, and to compare the succession of its members with the embryological development of the types now living in our seas. Ample material for making this comparison is fortunately at hand; it is material of a peculiar kind, not easily obtained, and which thus far has not greatly attracted the attention of zoölogists.

Interesting and important as are the earliest stages of embryonic development in the different classes of the animal kingdom, as bearing upon the history of the first appearance of any organ and its subsequent modifications, they throw but little light on the subject before us. What we need for our comparisons are the various stages of growth through which the young Sea-urchins of different families pass from the time they have practically become Sea-urchins until they have attained the stage which we now dignify with the name of species. Few embryologists have carried their investigations into the more extended field of the changes the embryo undergoes when it begins to be recognized as belonging to a special class, and when the knowledge of the specialist is absolutely needed to trace the bearing of the changes undergone, and to understand their full meaning. Fortunately the growth of the

young Echini has been traced in a sufficient number of families to enable me to draw the parallelism between these various stages of growth and the paleontological stages in a very different manner from what is possible in other groups of the animal kingdom, where we are overwhelmed with the number of species, as in the Insects or Mollusks, or where the paleontological or the embryological terms of comparison are wanting or very imperfect.

Beginning with the paleontological history of the regular Sea-urchins of the time of the Trias, when they constituted an unimportant group as compared with the Crenoids, we find the Echini of that time limited to representatives of two families. One of these, the genus *Cidaris*, has continued to exist, with slight modifications, up to the present time, and not less than one-tenth of all the known species of fossil Echini belong to this important genus, which in our tropical seas is still a prominent one. It is interesting here to note that in the *Cidaridæ* the modifications of the test are not striking, and the fossil genera appearing in the successive formations are distinguished by characters which often leave us in doubt as to the genus to which many species should be referred. In the genus *Rhabdocidaris*, which appears in the lower Jura, and which is mainly characterized by the extraordinary development of the radioles, we find the extreme of the variations of the spines in this family. From that time to the present day, the most striking differences have existed in the shape of the spines, not only of closely allied genera, but even in specimens of the same species; differences which in some of the species of today are as great as in older geological periods. The oldest *Cidaridæ* are remarkable for their narrow poriferous zones. It is only in the Jura that they widen somewhat; subsequently the pores become conjugated, and only later, during the Cretaceous period, do we find the first traces of any ornamentation of the test (*Temnocidaris*) so marked at the present day in the genus *Goniocidaris*. As far, then, as the *Cidaridæ* are concerned, the modifications which take place from their earliest appearance are restricted to slight changes in the poriferous zone and in the ornamentation of the test, accompanied with great variability in the shape of the primary radioles. We must except from this statement the genera *Diplocidaris* and *Tetracidaris*, to which I shall refer again. The representatives of the other Triassic family become extinct in the lower tertiary. The oldest genus, *Hemicidaris*, undoubtedly represents the earliest deviations from the true *Cidaris* type; modifications which affect not only the poriferous zone, but the test, the actinal and the abactinal systems, while from the extent of these minor changes we can trace out the gradual development of some of the characteristics in families of the regular Echini now living. The genus *Hemicidaris* may be considered as a *Cidaris* in which the poriferous zone is narrow and undulating, in which the granules of the ambulacral system have become minute tubercles in the upper portion of the zone and small primary tubercles in its actinal region, in which many of the interambulacral granules become small secondaries, in which the plates of the actinal system have become reduced in number, and the apical system has become a narrow ring, and finally in which the primary radioles no longer assume the fantastic shapes so common among the *Cidaridæ*.

We can trace in this genus the origin of the modifications of the poriferous zone, leading us, on the one side, through genera with merely undulating lines of pores to more or less distinct confluent arcs of pores, formed round the primary ambulacral tubercles, and, on the other, to the formation of open arcs of three or more pairs of pores. The first type culminates at the present day with the *Arbaciadæ*, the other with the *Diadematidæ*, *Triplechinidæ*, and *Echinometradæ*. This specialization very early takes place, for already in the lower Jura *Stomechinus* has assumed the principal characteristics of the *Triplechinidæ* of to-day.

Although in *Hemicidaris* the number of the coronal plates has increased as compared with the *Cidaridæ*, and while we find that in many genera, even of those of the present day, the number of the coronal plates is still comparatively small, yet, as a general rule, the more recent formations contain genera in which the increase in number of the interambulacral plates is accompanied by a corresponding decrease in the number of plates of the interambulacral area so characteristic thus far of the *Cidaridæ* and

Hemicidaridæ, a change also affecting the size of the primary ambulacral tubercles. This increase in the number of the coronal plates is likewise accompanied by the development of irregular secondary and miliary tubercles, and the disappearance in this group of the granular tuberculation, so important a character in the *Cidaridæ*. With the increase in the number of the interambulacral coronal plates, the *Pseudodiadematidæ* still retain prominent primary tubercles, recalling the earlier *Hemicidaridæ* and *Cidaridæ*, and, as in the *Cidaridæ* proper, the test is frequently ornamented by deep pits or by ridges formed by the junction of adjoining tubercles. The genital ring becomes narrower, and the tendency to the specialization of one of its plates, the madreporite, more and more marked.

With the appearance of *Stomechinus*, the *Echinidæ* proper already assume in the Jura the open arcs of pores, the large number of coronal interambulacral plates, the specialization of the secondary tubercles, and the large number of primary tubercles in each plate. With the appearance of *Sphærechinus* in the early Tertiary come in all the elements for the greater multiplication of the pairs of pores in the arcs of the poriferous zones, while the gigantic primary spines of some of the genera (*Heterocentrotus*), and the small number of primary tubercles are structural features which had completely disappeared in the group preceding the *Echinometradæ*, to which they appear most closely allied.

Going back again to the *Hemicidaridæ*, it requires but slight changes to pass from them to *Acrosalenia* and to the *Saleniæ* proper; the latter have continued to the present day, and have, like the *Cidaridæ*, retained almost unchanged the characters of the genera which preceded them, combined, however, with a few *Cidaridian* and *Echinid* features which date back to the Triassic period. We can thus trace the modifications which have taken place in the poriferous zone, the apical and actinal systems, the coronal plates, the ambulacral and interambulacral tubercles, as well as in the radioles, and in the most direct manner possible indicate the origin of the peculiar combination of structural features which we find at any geological horizon. On taking in succession the modifications undergone by the different parts of the test, we can trace each one singly, without the endless complication of combinations which any attempt to trace the whole of any special generic combination would imply.

Leaving out of the question for the moment the *Palæchinidæ*, we find no difficulty in tracing the history of the characters of the genera of the regular Echini which have existed from the time of the Trias and are now living, provided we take up each character independently. Nothing can be more direct than the gradual modification of the simple, barely undulating poriferous zone, made up of numerous ambulacral plates covered by granules, such as we find it among the *Cidaridæ* of Trias, first into the slightly undulating poriferous zone of the *Hemicidaridæ*, next into the indistinct arcs of pores of the *Pseudodiadematidæ*, then into the arcs with a limited number of pores of the *Triplechinidæ*, and finally to the polyporous arcs of the *Echinometradæ*. What can be more direct than the gradual modification to be traced in the development of the primary ambulacral tubercles, such as are characteristic of the *Echinidæ* of the present day, from their first appearance at the oral extremity of the ambulacral system of the *Hemicidaridæ*, and the increase in the number of primary interambulacral tubercles, accompanied by the growth of secondaries and miliaries, which we can trace in *Hemicidaridis*, *Acrosalenia*, and *Stomechinus*,—the increase in number of primary and secondary tubercles being accompanied by a reduction in the size of the radioles and a greater uniformity in their size and shape?

But while these modifications take place, the original structural feature may be retained in an allied group. Thus the *Cidaridæ* retain unchanged from the earliest time to the present day the few primary tubercles, the secondary granules, the simple poriferous zone, the imbricating actinal system, and the few coronal plates, with the large apical system and many-shaped radioles; while in the *Saleniæ* the primary internambulacral tubercles, the secondary granules, the radioles, the genital ring, are recognized features of the *Cidaridæ*, associated, however, with an *Echinid* actinal and anal system, *Hemicidarid* primary ambulacral

tubercles, and an Echinid poriferous zone. In the same way in the Diadematiadæ, the large primary internambulacral tubercles are Cidaridian features, while the structure of the ambulacral tubercles is Hemicidaridian. The existence of two kinds of spines is another Cidaridian feature, while the apical and actinal systems have become modified in the same direction as that of the Echinidæ. The more recent the genus, the greater is the difficulty of tracing in a direct manner the origin of any one structural feature, owing to the difficulty of disassociating structural elements characteristic of genera which may be derived from totally different sources. This is particularly the case with genera having a great geological age. Many of them, especially among the Spatangoids, show affinities with genera following them in time, to be explained at present only on the supposition that, when a structural feature has once made its appearance, it may reappear subsequently, apparently as a new creation, while in reality it is only its peculiar combination with structural features with which it had not before been associated (a new genus), which conceals in that instance the fact of its previous existence. A careful analysis, not only of the genera of the order, but sometimes of other orders which have preceded this combination in time, may often reveal the elements from which have been produced apparently unintelligible modifications.

There is, however, not one of the simple structural features in the few types of the Triassic and Liassic Echini from which we can so easily trace the origin of the structural features of all the subsequent Echinid genera, which is not also itself continued to the present day in some generic type of the present epoch, fully as well characterized as it was at the beginning. In fact, the very existence to-day of these early structural features seem to be as positive a proof of the unbroken systematic affinity between the Echini of our seas and those of the Trias, as the uninterrupted existence of the genus *Pygaster* or *Cidarid* from the Trias down to the present epoch, or of the connection of many of the genera of the Chalk with those of our epoch (*Salenia*, *Cyphosoma*, *Psammechinus*, etc.).

Passing to the Clypeastridæ, we find there as among the Desmosticha the earliest type, *Pygaster*, has existed from the Trias to the present time; and that, while we can readily reconstruct, on embryological grounds, the modifications the earliest Desmosticha-like Echini should undergo in order to assume the structural features of *Pygaster*, yet the early periods in which the precursors of the Echinoconidæ and Clypeastridæ are found have thus far not produced the genera in which these modifications actually take place. But, starting from *Pygaster*, we naturally pass to *Holactypus*, to *Discoidea*, to *Conoclypus*, on the one side, while on the other, from *Holactypus* to *Echinocyamus*, *Sismondia*, *Fibularia*, and *Mortonia*, we have the natural sequence of the characters of the existing Echinanthidæ, *Laganidæ*, and *Scutellidæ*, the greater number of which are characteristic of the present epoch. If we were to take in turn the changes undergone in the arrangement of the plates of the test, as we pass from *Pygaster* to *Holactypus*, to *Echinocyamus*, and *Echinanthidæ*, we should have in the genera which follow each other in the paleontological record an unbroken series showing exactly what these modifications have been. In the same way, the modifications of the abactinal and anal systems, and those of the poriferous zone, can equally well be followed to *Echinocyamus*, and thence to the Clypeastridæ; while a similar sequence in the modifications of these structural features can be followed from *Mortonia* to the *Scutellidæ* of the present period.

Passing finally to the Petalosticha, we find no difficulty in tracing theoretically the modifications which our early Echinoconidæ of the Lias should primarily undergo previous to the appearance of *Galeropygus*. The similarity of the early Cassiduloid and Echinoneoid types points to the same systematic affinity, and perhaps even to a direct and not very distant relationship with the Palæchinidæ. For if we analyze the Echinothuridæ of the present day, we find in genera like *Phormosoma* many structural features, such as the shape of the test the character of the spines, the structure of the apical system, that of the poriferous zone, indicative of possible modifications in the direction of *Pygaster* or of *Galeropygus*, which have as yet not been taken into account.

Adopting for the Petalosticha the same method of tracing the modifications of single structural features in their paleontological succession, we trace the comparatively little modified paleontological history of the Echinoneidæ of the present day from the *Pyrina* of the lower Jura. This, in its turn, has been preceded by *Hyboclypus* and *Galeropygus*, while the Echinolampadæ of the present day date back, with but trifling modifications, to the Echinobrissus of the Lias, itself preceded by *Clypeus*; and they have been subject only to slight generic changes since that time, Echinobrissus being still extant, while such closely allied genera as *Catopygus* and *Cassidulus* of the earlier Cretaceous are still represented at the present day; the modifications taking place in the actinal system, in the ambulacral zones of the Echinoconidæ and of the Echinolampadæ showing the closest possible systematic affinity in these families. Starting again from *Hyboclypus*, with its elongate apical system, we naturally pass to *Collyrites* and the strange *Dysasteridæ* forms which, in their turn, are closely allied to the *Holasteridæ*. From *Holaster* on the one side, and from *Toxaster* on the other, we find an unbroken sequence of structural characters uniting the successive genera of *Holasteridæ*, such as *Cardiaster*, *Offaster*, *Stenonia*, *Ananchytes*, and *Asterostoma*, with *Paleopneustes*, *Homolampas* and the *Pourtalesidæ* of the present day, while from the genera of the *Toxasteridæ* we naturally pass to the Cretaceous *Hemiaster*; in the genus and the subsequent *Micraster* we find all the elements necessary for the modifications which appear in the Spatanginæ from the time of the Chalk to the present day. These modifications result in genera in which we trace the development of the fascioles, of the actinal, anal, and abactinal plastrons, of the beak, the formation of the petaloid ambulacra, first flush with the test, and little by little changed into marsupial pouches, the growth of the anterior groove and the manifold modifications of the ambulacral system in *Spatangus*, *Agassizia*, and *Echinocardium*, often recalling in some of its features structural characters of families which have preceded this in time.

Apparently in striking contrast with the Echini of the secondary period and those which have succeeded them stand the Paleozoic Echini; but when we have examined the embryology of Echini, we shall be better prepared to understand their structure and the affinities of the Palæchinidæ with the Echini of the present day and their immediate predecessors.

Taking up now the embryological development of the several families which will form the basis of our comparisons, beginning with the Cidaridæ, we find that in the earliest stages they very soon assume the characters of the adult, the changes being limited to the development of the abactinal system, the increase in number of the coronal plates, and the modifications of the proportionally gigantic primary radioles.

In the Diadematiadæ the changes undergone by the young are limited to the gradual transformation of the embryonic spines into those which characterize the family, to the changes of the vertical row of pores in the ambulacral area into arcs of three or four pairs of pores, and to the specialization of the actinal and abactinal systems.

In the Arbaciadæ the young stages are remarkable for the prominent sculpture of the test, for the flattened spines, for their simple poriferous zone, for their actinal system, and for their genital ring. The anal plates appear before the genital ring.

In the Echinometradæ the young thus far observed are characterized by the small number of their primary tubercles, the large size of the spines, the simple vertical row of pores, the closing of the anal ring by a single plate, and the turban-shaped outline of the test. Little by little, the test loses with increasing age this Cidarid-like character; it reminds us, from the increase in the number of its plates, more of *Hemicidarid*; then, with their still greater increase, of the *Pseudodiadematiadæ*; and, finally, of the *Echinometradæ* proper. The spines, following *pari passu* the changes of the test, lose little by little their fantastic embryonic, or rather Cidarid-like appearance, and become more solid and shorter, till they finally assume the delicately fluted structure characteristic of the Echinometradæ. The vertical poriferous zone is first changed into a series of connected vertical arcs, which become disjointed, and form, with increas-

ing age, the independent arcs of pores, composed of three or more pair of pores, of the Echinometradæ.

In the Echinidæ proper we find in the young stages the same unbroken vertical line of pores, which gradually becomes changed to the characteristic generic types. We find, as in the Echinometradæ, an anal system closed with a single plate, and an abactinal system separating in somewhat more advanced stages from the coronal plates of the test. This is as yet made up of a comparatively small number of plates, carrying but few large primary tubercles, with fantastically shaped spines entirely out of proportion to the test, but which, little by little, with the increase of the number of coronal plates, the addition of primary tubercles, and their proportional decrease in size, assume more and more the structure of the genus to which the young belongs. The original anal plate is gradually lost sight of from the increase in number of the plates covering the anal system, and it is only among the Temnopleuridæ that this anal plate remains more or less prominent in the adult. In the Salinidæ, of which we know as yet nothing of the development, this embryonic plate remains permanently a prominent structural feature of the apical system.¹

Among the Clypeastroids the changes of form they undergo during growth are most instructive. We have in the young Fibularinæ an ovoid test, a small number of coronal plates surmounted by few and large primary tubercles, supporting proportionally equally large primary radioles, simple rectilinear poriferous zones, no petaloid ambulacra,—in fact, scarcely one of the features we are accustomed to associate with the Clypeastroids is as yet prominently developed. But rapidly, with increasing size, the number of primary tubercles increases, the spines lose their disproportionate size, the pores of the abactinal region become crowded and elongate, and a rudimentary petal is formed. The test becomes more flattened, the coronal plates increase in number, and it would be impossible to recognize in the young Echinocyamus, for instance, the adult of the Cidaridæ-like or Echinometra-like stages of the Sea-urchin, had we not traced them step by step. Most interesting, also, is it to follow the migrations of the anal system, which, to a certain extent, may be said to retain the embryonic features of the earlier stages of all Echinoderm embryos, in being placed in more or less close proximity to the actinostome. What has taken place in the growth of the young Echinocyamus is practically repeated for all the families of Clypeastroids: a young Echinarachnius, or Mellita, or Encope, or a Clypeaster proper, resembles at first more an Echinometra than a Clypeastroid; they all have simple poriferous zones and spines and tubercles out of all proportion to the size of the test.²

When we come to the development of the Spatangoids, we find their younger stages also differing greatly from the adult. Among the Nucleolidæ, for instance, the young stages have as yet no petals, but only simple rectilinear poriferous zones. They are elliptical with a high test, with a single large primary tubercle for each plate, and a simple elliptical actinostome, without any trace of the typical bourrelets and phyllodes so characteristic of this family. Very early, however, this condition of things is changed, the test soon becomes more flattened, the petals begin to form as they do in the Clypeastroids, and we can soon trace the rudiments of the peculiar bourrelets characteristic of the family, accompanied by a rapid increase in the number of tubercles and in that of the coronal plates.

Among the Spatangidæ some are remarkable in their adult condition for their labiate actinostome, for the great development of the petals, for the presence of fascioles surrounding certain definite areas, for the small size of the tubercles, the general uniformity in the spines of the test, and the specialization of their anterior and posterior regions. On examining the young stages of this group of Spatangoids, not one of these structural features is as yet developed. The actinostome is simple, the poriferous zone has the same

simple structure from the actinostome to the apex, the primary tubercles are large, few in number, surrounded by spines which would more readily pass as the spines of Cidaridæ than of Spatangoids. The fascioles are either very indistinctly indicated, or else the special lines have not as yet made their appearance; the ambulacral suckers of the anterior zone are as large and prominent as those of the young stages of any of the regular Echini. It is only little by little, with advancing age, that we begin to see signs of the specialization of the anterior and posterior parts of the test, that we find the characteristic anal or lateral fascioles making their appearance, only with increasing size that the spines lose their Cidaridæ-like appearance, that the petals begin to be formed, and that the simple actinostome develops a prominent posterior lip. In the genus Hemiaster, the young stages are especially interesting, as long before the appearance of the petals, while the poriferous zone is still simple, the total separation of the bivium and of the trivium of the ambulacral system, so characteristic of the earliest Spatangoids (the Dysasteridæ), is very apparent.³

From this rapid sketch of the changes of growth in the principal families of the recent Echini we can now indicate the transformations of a more general character through which the groups as a whole pass.

In the first place, while still in the Pluteus all the young Echini are remarkable for the small number of coronal plates, and for the absence of any separation between the actinal and abactinal systems and the test proper. They all further agree in the large size of the primary spines of the test, whether it be the young of a Cidaridæ, an Arbacia, an Echinus, a Clypeaster, or a Spatangoid. They all in their youngest stages have simple vertical ambulacral zones; beyond this, we find as changes characteristic of some of the Desmostichæ, the specialization of the actinal system from the coronal plates, the formation of an anal system, the rapid increase in the number of the coronal plates, with a corresponding increase in the number of the spines and a proportional reduction of their size, the formation of an abactinal ring, and the change of the simple vertical poriferous zone into one composed of independent arcs.

In the Spatangoids and Clypeastroids we find common to both groups the shifting of the anal system to its definite place, the modifications of the abactinal part of the simple ambulacral system in order to become petaloid, and the gradual change of the elliptical ovoid test of the young to the characteristic generic test, accompanied by the rapid increase in the number of the primary tubercles and spines. Finally limited to the Spatangoids are the changes they undergo in the transformation of the simple actinostome to a labiate one, the specialization of the anterior and posterior parts of the test, and the definite formation of the fascioles.

Comparing this embryonic development with the paleontological one, we find a remarkable similarity in both, and in a general way there seems to be a parallelism in the appearance of the fossil genera and the successive stages of the development of the Echini as we have traced it.

We find that the earlier regular Echini all have more or less a Cidaridæ-like look,—that is, they are Echini with few coronal plates; large primary tubercles, with radioles of a corresponding size; that it is only somewhat later that the Diademopsidæ make their appearance, which, in their turn, correspond within certain limits to the modifications we have traced in the growth of the young Diadematiidæ and Arbaciadæ. The separation of the actinal system from the coronal plates has been effected. The poriferous zone has either become undulating, or forms somewhat indefinite open arcs; we find in all the genera of this group a larger number of coronal plates, more numerous primaries, the granules of the Cidaridæ replaced by secondaries and miliaries, and traces of a Hemicidaridæ-like stage in the size of the actinal ambulacral tubercles.

Comparing in the same way the paleontological development of the Echinidæ proper, we find that, on the whole, they agree well with the changes of growth we can still follow to-day in their representatives, and that, as we approach nearer the present epoch, the fossil genera more and more assume the structural features which we find de-

¹ The young of the following genera have served as a basis for the preceding analysis of the embryonic stages of the Desmostichæ: Cidaridæ, Dorocidaridæ, Gonocidaridæ, Arbacia, Porocidaridæ, Strongylocentrotus, Echinometra, Echinus, Tæxopneustes, Hipponoë, Temnopleurus, Temnechinus, and Trigonocidaridæ.

² Among the Clypeastroids I have examined the young of Echinocyamus, Fibularia, Mellita, Laganum, Echinarachnius, Encope, Clypeaster, and Echinanthus.

³ For this sketch of the embryology of the Petalostichæ I have examined the young of Echinolampus, Echinoneus, Echinocardium, Brissopsis, Agassizia, Spatangus, Brissus, and Hemiaster.

veloped last among the Echinidæ of the present day. Very much in the same manner as a young Echinus develops, they lose, little by little, first their Cidaridian affinities, which become more and more indefinite, next their Didemmatidian affinities, if I may so call the young stages to which they are most closely allied, and, finally, with the increase in the number of the coronal plates, the great numerical development of the primary tubercles and spines, and that of the secondaries and miliaries which we can trace in the fossil Echini of the Tertiaries, we pass insensibly into the generic types characteristic of the present day.

Although we know nothing of the embryology of the Salenidæ, yet, like the Cidaridæ, they have in a great measure remained a persistent type, the modifications of the group being all in the same direction as those noticed in the other Desmosticha; a greater number of coronal plates; the development of secondaries and miliaries combined with a specialization of the actinal system not found in the Cidaridæ.

An examination of the succession of the Echinoconidæ shows but little modification from the earliest types; the changes, however, are similar to those undergone by the Clypeastroids and Petalosticha, though they do not extend to modifications of the poriferous zone, but are mainly changes in the actinostome and in the tuberculation. In fact, the group of Echinoconidæ seems to hold somewhat the same relation to the Clypeastroids which the Salenidæ hold to the Cidaridæ, and the earliest genus of the group (Pygaster) has remained, like Cidaris, a persistent type to the present day.

The earliest Clypeastroids are all forms which resemble the Fibularina and the genera following Echinocyamus and Fibularia; they are mainly characterized by the same changes which an Echinarachnius or a Mellita, for instance, undergoes as it passes from its Echinocyamus stage to the Laganum or Encope stage. The comparison is somewhat more complicated when we come to the Spatangoids. The comparison of the succession of genera in the different families, as traced in the Desmosticha and Clypeastroids, is made difficult from the persistency of the types preceding the Echinoneidæ and the Ananchytidæ, which have remained without important modifications from the time of the lower Cretaceous; previous to that time the modifications of the Cassidulidæ are found to agree with the changes which have been observed in the growth of Echinolampas. The early genera, like Pygurus, have many of the characteristics of the test of the young Echinolampas. The development of prominent bourrelets and of the floscelle and petals goes on side by side with that of genera in which the modification of the actinostome, of the test, and of the petals is far less rapid, one group retaining the Echinoneus features, the other culminating in the Echinolampas of the present day, and having likewise a persistent type, Echinobrissus, which has remained with its main structural features unchanged from the Jura to the present day. That is, we find genera of the Cassidulidæ which recall the early Echinoneus stage of Echinolampas, next the Caratomus stage, after which the floscelle, bourrelets, and petals of the group become more prominent features of the succeeding genera. Accompanying the persistent type Echinobrissus, genera appear in which either the bourrelets or petals have undergone modifications more extensive than those of the same parts in the genera of the Echinoneus or Caratomus type.

The earliest Spatangoids belong to the Dysasteridæ, apparently an aberrant group, but which, from the history of the young Hemiaster, we now know to be a strictly embryonic type, which, while it thus has affinities with the true Spatangoids, still retains features of the Cassidulidæ in the mode of development of the actinostome and of the petals, as well as of the anal system. The genera following this group, Holaster and Toxaster, can be well compared, the one to the young stages of Spatangus proper before the appearance of the petals, when the ambulacra are flush with the test, and when its test is more or less ovoid, the other to a somewhat more advanced stage, when the petals have made their appearance as semi-petals. In both cases the actinostome has the simple structure characteristic of all the young Spatangoids. The changes we notice in the genera which follow them lead in the one case through very slight modifications of the abactinal system, of the anterior

and posterior extremities of the test, to the Ananchytid-like Spatangoids of the present day, the Pourtalesidæ, the genus Holaster itself persisting till well into the middle of the Tertiary period; while on the other side we readily recognize in the Spatanginæ which follow Toxaster (a persistent type which has continued as Palæostoma to the present day) the genera which correspond to the young stages of such Spatangoids as Spatangus and Brissopsis of the present day, genera which, on the one hand, lead from Hemiaster (itself still represented in the present epoch), through stages such as Cyclaster, Peripneustes, Brissus, and Schizaster, and, on the other, through Micraster and the like, to the Spatangoids, in which the development of the anal plastron and fasciole performs an important part, while in the former group the development of the peripetalus fasciole and of the lateral fasciole can be followed. None of the genera of Petalosticha belonging to the other groups develop any fasciole in the sense of circumscribing a limited area of the test.

The comparison of the genera of Echini which have appeared since the Lias with the young stages of growth of the principal families of Echini, shows a most striking coincidence amounting almost to identity between the successive fossil genera and the various stages of growth. This identity can, however, not be traced exactly in the way in which it has usually been understood, while there undoubtedly exists in the genera which have appeared one after the other a gradual increase in certain families in the number of forms, and a constant approach in each succeeding formation, in the structure of the genera, to those of the present day. It is only in the accordance between some special points of structure of these genera and the young stages of the Echini of the present day that we can trace an agreement which, as we go further back in time, becomes more and more limited. We are either compelled to seek for the origin of many structural features in types of which we have no record, or else we must attempt to find them existing potentially in groups where we had as yet not succeeded in tracing them. The parallelism we have traced does not extend to the structure as a whole. What we find is the appearance among the fossil genera of certain structural features giving to the particular stages we are comparing their characteristic aspect. Thus, in the succession of the fossil genera, when a structural feature has once made its appearance, it may either remain as a persistent structure, or it may become gradually modified in the succeeding genera of the same family, or it may appear in another family, associated with other more marked structural features which completely overshadow it. Take, for instance, among the Desmosticha, the modifications of the poriferous zone of the actinal and abactinal systems of the coronal plates, of the ambulacral and interambulacral systems, the changes in the relative proportion of the primary-tubercles, and the development of the secondaries. These are all structural features which are modified independently one of the other; we may find simultaneous development of these features in parallel lines, but a very different degree of development of any special feature in separate families.

This is as plainly shown in the embryological as in the paleontological development. In the Cidaridæ there is the minimum of specialization in these structural features. In the Diademopsidæ there is a greater range in the diversity of the structure of the poriferous zone and of the coronal plates, as well as of the actinal system. There is a still greater range among the Echinidæ, while among the Salenidæ the modifications, as compared to those of the Echinidæ and Diademopsidæ, are somewhat limited again, being restricted as far as relates to the poriferous zone and coronal plates, but specialized as far as the actinal system is concerned, and specially important with reference to the structure of the apical system. The special lines in which these modifications take place produce, of course, all possible combinations, yet they give us the key to the sudden appearance, as it were, of structural features of which the relationship must be sought in very distantly related groups. It is to this specialty in the paleontological development that we must trace, for instance, the Cidarid affinities of the Salenidæ, their papillæ, the existence of few large primary interambulacral tubercles, the structure of their apical system, and their large genital plates; while it is to

their affinities with the Hemicidaridæ that we must refer the presence of the few larger primary ambulacral tubercles at the base of the ambulacral area, and by their Diademopod and Echinidian affinities that we explain the indented imbricated actinal system with the presence of a few genuine miliaries. But all the structural features which characterize the earliest types of the Desmosticha can in reality be traced, only in a somewhat rudimentary form, even in the Cidaridæ. The slight undulation of the closely packed, nearly vertical poriferous zone is the forerunner of the poriferous zone first separated into vertical arcs and then into independent arcs. The limitation in the number of the rows of granules in the ambulacral zone, and their increase in size, is the first trace of the appearance of the somewhat larger primary ambulacral tubercles of the Hemicidaridæ and Salenidæ. The existence of the smooth cylindrical spines of the abactinal region of the test naturally leads to similar spines covering the whole test in the other families in the Desmosticha. The difference existing in the plates covering the actinal system from those of the coronal plates leads to the great distinction between the structure of the actinal system and of the coronal plates in some of the Echinidæ.

Passing to the Clypeastridæ and Petalosticha, we trace a parallelism of the same kind, and readily in the successive genera of fossil Clypeastroids, but often in widely separated genera, the precise modifications which the poriferous zone has undergone as it first becomes known to us in Echinocyamus and Fibularia, and as we find it in the most complicated petaloid stage of the Clypeastroids of the present day. We readily trace the changes the test undergoes from its comparatively ovoid and swollen shape to assume first that of the less gibbous forms, next that of the Laganidæ, and finally of the flat Scutellidæ; while we trace in the Echinanthidæ the persistent structural features of some of the earliest Clypeastroids, together with an excessive modification of the poriferous zone. Likewise for the Echinocnidæ we trace mainly the slight modifications of the poriferous zone and of the coronal plates, and finally, when we come to the Spatangidæ we find no difficulty in tracing from the most Desmostichoid of the Spatangoid genera, the modifications of a test in which the ambulacral and interambulacral areas are made up of plates of nearly uniform size, in which the anterior and posterior extremities are barely specialized, to the most typical of the Ananchytidæ, in which the anterior and posterior extremities have developed the most opposite and extraordinary structural features. In a similar way we can trace among the fossil genera of different families the gradual development of the actinal plastron from its very earliest appearance as a modification of the posterior interambulacral area of the actinal side, or the growth of the posterior beak into an anal snout, the successive changes of the anal groove, the formation of the actinal labium, or the development of the bourrelets and phyllodes from a simple circular actinostome, the gradual deepening of the slight anterior groove of some early Spatangoid to form the deeply sunken actinal groove. Equally well we can trace the modifications of the ambulacral system as it passes from the simple poriferous zones of the earlier Spatangoids to genera in which the petaliferous portion makes its appearance, and finally becomes the specialized structure of our recent Spatangoid genera, such as Schizaster, Moira, and the like. Finally we can trace, to a certain extent, the development of the fascioles on one side from genera like Hemiaster, in which the peripetalous fasciole is prominent, to genera like Brissopsis, Brissus, and the like, of the present day; on the other, perhaps, or in both combined, the formation of a lateral and anal fasciole from genera like Micraster in Spatangus and Agassizia. Thus we must, on the same theory of the independent modifications of special structural features, trace the many and complicated affinities which so constantly strike us in making comparative studies, and which render it impossible for us to express the manifold affinities we notice, without taking up separately each special structure. Any attempt to take up a combination of characters, or a system of combinations, is sure to lead us to indefinite problems far beyond our power to grasp.

In the oldest fossil Clypeastroids and Petalosticha, as well as in the Desmosticha, we also find the potential expression of the greater number of the modifications subse-

quently carried out in genera of later date. The semipetaloid structure of some of the earlier genera of Spatangoids, the slight modifications of some of the plates of the actinal side near the actinostome, are the precursors, the one of the highly complicated petaloid ambulacra of the recent Spatangoids, the other of the actinal plastron, leading as it does also to the important differences subsequently developed in the anterior and posterior extremities of the test, as well as to the modifications which lead to the existence of a highly labiate actinostome. The appearance of a few miliaries near the actinostome constitutes the first rudimentary bourrelets.

Going back now to the Palæchinidæ, the earliest representatives of the Echini in paleozoic times, without any attempt to trace the descent of any special type from them, we may perhaps find some clew to the probable modifications of their principal structural features preparatory to their gradual disappearance. In the structure of the coronal plates, the specialization of the actinal and abactinal systems, the conditions of the ambulacral system, we must compare them to stages in the embryonic development of our recent Echini with which we find no analogues in the fossil Echini of the Lias and the subsequent formations. In order to make our parallelism, we must go back to a stage in the embryonic history of the young Echini in which the distinction to be made between the ambulacral and interambulacral systems is very indefinite, in which the apical system is, it is true, specialized, but in which the actinal system remains practically a part of the coronal system. But here the comparison ceases, and, although we can trace in the paleontological development of such types as Archæocidaris or Bothriocidaris modifications which would lead us without great difficulty, on the one side to the Cidaridæ, and on the other to the Echinothuriæ and Diademantidæ of the present day, we cannot fail to see most definite indications in some of the structural features of the Palæchinidæ of characteristics which we have been accustomed to associate with higher groups. The minute tuberculation, for instance, of the Clypeastroids and Spatangoids, already existing in the Melonitidæ, the genital ring, and anal system, are quite as much Echinid as Cidarid. The polyporous genera of the group represent to a certain extent the polypori of the regular Echini, and the lapping of the actinal plates of the Cidaridæ and of the coronal plates in some of the Diademantidæ, as well as the existence of such genera as Tetracidaris, of four interambulacral plates in Astropyga, and of a large number of ambulacral plates in some of the recent Echinometradæ, all these are Palæchinid characters which we can explain on the theory of the independent development of the structural features of which they are modifications. We should, however, remember, that the existence of a large number of coronal plates, especially interambulacral plates, in the Palæchinidæ, is a mere vegetative character, which they hold in common with all the Crinoids—a character which is reduced to a minimum among the Holothurians, and still persists in full force among the Pentacrinini of the present day, as well as the Astrophytidæ and Echinidæ.

It would lead me too far to institute the same comparison between the embryonic stages of the different orders of Echinoderms and their earliest fossil representatives. We may, however, in a very general way, state that we know the earliest embryonic stages of the orders of Echinoderms of to-day, which, with the exception of the Blastoidæ and Cystideans, are identical with the fossil orders, and that as far as we know they all begin at a stage where it would be impossible to distinguish a Sea-urchin from a Star-fish, or an Ophiuran, or a Crinoid, or an Holothurian—a stage in which the test, calyx, abactinal and ambulacral systems are reduced to a minimum. From this identical origin there is developed at the present day, in a comparatively short period of time, either a Star-fish, a Sea-urchin, or a Crinoid; and if we have been able successfully to compare, in the development of typical structures, the embryonic stages of the young Echini with their development in the fossil genera, we may fairly assume that the same process is applicable when instituting the comparison within the different limits of the orders, but with the same restrictions. That is, if we wish to form some idea of the probable course of transformations which the earliest Echinoderms have undergone to lead us to those of the present day, we are

justified in seeking for our earliest representatives of the orders such Echinoderms as resemble the early stages of our embryos, and in following, for them as for the Echini, the modifications of typical structures. These we shall have every reason to expect to find repeated in the fossils of later periods, and, going back a step further, we may perhaps get an indefinite glimpse of that first Echinodermal stage which should combine the structural features common to all the earliest stages of our Echinoderm embryos.

And yet, among the fossil Echinoderms of the oldest periods, we have not as yet discovered this earliest type from which we would derive either the Star-fishes, Ophiurans, Sea-urchins, or Holothurians. With the exception of the latter, which we can leave out of the question at present, we find all the orders of Echinoderms appearing at the same time. But while this is the case, one of the groups attained in these earliest days a prominence which it gradually loses with the corresponding development of the Star-fishes, Ophiurans, and Sea-urchins, it has steadily declined in importance; it is a type of Crinoids, the Cystideans which culminated during Paleozoic times, and completely disappeared long before the present day. If we compare the early types of Cystideans to the typical embryonic Echinodermal type of the present day, we find they have a general resemblance, and that the Cystideans and Blastoids represent among the fossil Echinoderms the nearest approach we have yet discovered to this imaginary prototype of Echinoderms.

This may not seem a very satisfactory result to have attained. It certainly has been shown to be an impossibility to trace in the paleontological succession of the Echini anything like a sequence of genera. No direct filiation can be shown to exist, and yet the very existence of persistent types, not only among Echinoderms, but in every group of marine animals, genera which have continued to exist without interruption from the earliest epochs at which they occur to the present day, would prove conclusively that at any rate some groups among the marine animals of the present day are the direct descendants of those of the earliest geological periods. When we come to types which have not continued as long, but yet which have extended through two or three great periods, we must likewise accord to their latest representatives a direct descent from the older. The very fact that the ocean basins date back to the earliest geological periods, and have afforded to the marine animals the conditions most favorable to an unbroken continuity under slightly varying circumstances, probably accounts for the great range in time during which many genera of Echini have existed. If we examine the interlacing in the succession of the genera characteristic of later geological epochs, we find it an impossibility to deny their continuity from the time of the Lias to the present day. The *Cidaris* of the Lias and the *Rhabdocidaris* of the Jura are the ancestors of the *Cidaris* of to-day. The *Salenia* of the lower Chalk are those of the *Salenia* of to-day. *Acrosalenia* extends from the Lias to the lower Cretaceous, with a number of recent genera, which begin at the Eocene. The *Pygaster* of to-day dates back to the Lias; *Echinocyamus* and *Fibularia* commence with the Chalk. *Pyrina* extends from the lower Jura through the Eocene. The *Echinobrissus* of to-day dates back to the Jura. *Holaster* lived from the lower Chalk to the Miocene, and the *Hemiaster* of to-day cannot be distinguished from the *Hemiaster* of the lower Cretaceous.

Such descent we can trace, and trace as confidently as we trace a part of the population of North America of to-day as the descendants of some portion of the population of the beginning of this century. But we can go no further with confidence, and bold indeed would he be who would attempt even in a single State to trace the genealogy of the inhabitants from those of ten years before. We had better acknowledge our inability to go beyond a certain point; anything beyond the general parallelism I have attempted to trace, which in no way invalidates the other proposition, we must recognize as hopeless.

But in spite of the limits which have been assigned to this general parallelism, it still remains an all-essential factor in elucidating the history of paleontological development, and its importance has but recently been fully appreciated. For, while the fossil remains may give us a strong presumptive evidence of the gradual passage of one type to another, we can only imagine this modification to take place by a process similar to that which brings about the modifications due

to different stages of growth,—the former taking place in what may practically be considered as infinite time when compared to the short life history which has given us as it were a *résumé* of the paleontological development. We may well pause to reflect that in the two modes of development we find the same periods of rapid modifications occurring at certain stages of growth or of historic development, repeating in a different direction the same phases. Does it then pass the limits of analogy to assume that the changes we see taking place under our own eyes in a comparatively short space of time,—changes which extend from stages representing perhaps the original type of the group to their most complicated structures,—may, perhaps, in the larger field of paleontological development, not have required the infinite time we are in the habit of asking for them?

Paleontologists have not been slow in following out the suggestive track, and those who have been anatomists and embryologists besides have not only entered into most interesting speculations regarding the origin of certain groups, but they have carried on the process still further, and have given us genealogical trees where we may, in the twigs and branches and main limbs and trunk, trace the complete filiation of a group as we know it to-day, and as it must theoretically have existed at various times to its very beginning. While we cannot but admire the boldness and ingenuity of these speculations upon genetic connection so recklessly launched during the last fifteen years, we find that with but few exceptions there is little to recommend in reconstructions which shoot so wide of the facts as far as they are known, and seem so readily to ignore them. The moment we leave out of sight the actual succession of the fossils and the ascertainable facts of post-embryonic development, to reconstruct our genealogy, we are building in the air. Ordinarily, the twigs of any genealogical tree have only a semblance of truth; they lead us to branchlets having but a slight trace of probability, to branches where the imagination plays an important part, to main limbs where it is finally allowed full play, in order to solve with the trunk, to the satisfaction of the writer at least, the riddle of the origin of the group. It seems hardly credible that a school which boasts for its very creed a belief in nothing which is not warranted by common sense should descend to such trifling.

The time for genealogical trees is passed; its futility can, perhaps, best be shown by a simple calculation, which will point out at a glance what these scientific arboriculturists are attempting. Let us take, for instance, the ten most characteristic features of Echini. The number of possible combinations which can be produced from them is so great that it would take no less than twenty years, at the rate of one new combination a minute for ten hours a day, to pass them in review. Remembering now that each one of these points of structure is itself undergoing constant modifications, we may get some idea of the nature of the problem we are attempting to solve, when seeking to trace the genealogy as understood by the makers of genealogical trees. On the other hand, in spite of the millions of possible combinations which these ten characters may assume when affecting not simply a single combination, but all the combinations which might arise from their extending over several hundred species, we yet find that the combinations which actually exist—those which leave their traces as fossils—fall immensely short of the possible number. We have, as I have stated, not more than twenty-three hundred species actually representing for the Echini the results of these endless combinations. Is it astonishing, therefore, that we should fail to discover the sequence of the genera, even if the genera, as is so often the case, represent, as it were, fixed embryonic stages of some Sea-urchin of the present day? In fact, does not the very history of the fossils themselves show that we cannot expect this? Each fossil species, during its development, must have passed through stages analogous to those gone through by the Echini of the present day. Each one of these stages at every moment represents one of the possible combinations, and those which are actually preserved correspond only to the particular period and the special combination which any Sea-urchin has reached. These stages are the true missing links, which we can no more expect to find preserved than we can expect to find a record of the actual embryonic development of the species of the present day without direct

observation at the time. The actual number of species in any one group must always fall far short of the possible number, and for this reason it is out of the question for us to attempt the solution of the problem of derivation, or to hope for any solution beyond one within the most indefinite limits of correctness. If, when we take one of the most limited of the groups of the animal kingdom, we find ourselves engaged in a hopeless task, what must be the prospect should we attack the problem of other classes or groups of the animal kingdom, where the species run into the thousands, while they number only tens in the case we have attempted to follow out? Shall we say "ignorabimus," or "impavidi progrediamur" and valiantly chase a phantom we can never hope to seize?

CHEMISTRY AS AN ART, AND CHEMISTRY AS A SCIENCE.

BY PROFESSOR J. M. ORDWAY.

Professor J. M. Ordway, of Boston, spoke of "Chemistry as an Art, and Chemistry as a Science," comparing both, and pointing out some recent lines of advancement. The past year, he said, has been one of laborious activity in chemistry, but it has not been marked by any epoch-making discoveries. Meyer's recent apparent resolution of the chlorine molecule has not, indeed, been verified by the carefully devised experiments of Crafts, but the latter does seemingly confirm the change of iodine by intense heat. The years 1879 and 1880 will rank hereafter as years in which Meyer found means to throw new light on the nature of the haloids. Twenty-four years ago Perkins sought for artificial quinine, and found instead a better than royal purple. Then, by various hands and in rapid succession, red and yellow and black and brown and blue dyes were brought out from what proved to be something more than aniline. Now the novelty is past, and the announcement of a new dye hardly creates a ripple of excitement. The twelve-year-old synthesis of alizarine has given us colors purer, brighter, faster and cheaper than those of the obsolescent madder. Of late, wool has been provided for, and the extinction of cochineal plantations is threatened by reds of surpassing brilliancy, durability and ease of application. Baeyer has recently effected the synthesis of indigo, and tropical indigo fields may in time share the fate of the madder farms of France and Turkey. But indigo itself will not continue to satisfy our demand. We have become accustomed to hues of a delicacy and richness that no one dared to dream of twenty-five years ago. The æsthetic taste of this generation has been too much pampered; and dyers will soon call for something uniting the brilliancy of the aniline blues with the fixedness of indigo, and its adaptedness to wool and cotton. And Germany which has done the most in studying out these extraordinary colored compounds, now furnished the most of the industrial fruits of seemingly unpractical researches. Investigation costs, investigation pays; in more senses than one our science "opens wide her everdaring gates on golden hinges turning."

The passing years are bringing to light new elementary bodies, and new metals are becoming like new asteroids, of too little mass to influence the orbits of other planets, and too much out of sight to interest many. Within five years fourteen new metals have called for recognition; and in 1879 alone chemists have claimed the discovery of six. Of new alloys, manganian copper is worthy of regard,

since it may in a measure play the part for copper that *spiegeleisen* does for steel.

In 1620 Bacon published the second part of his "Novum Organum," wherein he pointed out the way to appeal to nature by experiment, instead of deriving all science from the teaching of the ancients. But his methods had little immediate influence on the science of the time. He relied on induction; and induction alone simply strings together dry bones. That perception of general principles which makes science comes not altogether from the mere collation of facts. We need something more than eyes to see.

The great chemist of two hundred and fifty years ago was Van Helmont. To him we owe the word gas, which he derived not from *geist*, but from chaos, as representing the original form of matter. When our forefathers were laying the foundations of this nation alchemy was in its dotage, and chemistry took its rise in a dim knowledge of the gases. The evolution of chemistry as a science was threefold. First, the study of the gases, then the study of heat, then the study of combining weights. Consider how much of what we now know depends on the gases that Cavendish, Black, Scheele and Priestley revealed. The study of combustion, respiration, vegetable growth, organic decay, geological transformation and hygiene involves the study of carbon dioxide. Carbon monoxide reduces the metals, and plays a part in the Bessemer process for making steel. The fuel of the future is to be coal resolved into a chaos of carbonic oxide and hydrogen. At the end of the last century Murdoch found a use for coal gas, and in its train came a host of secondary products having a marvellous effect on science and industry. A test came into chemistry when Becher attempted to explain combustion. Vulcan of old made as good iron as the blacksmith requires to-day. As for quantity, Vulcan with all his Cyclops and the fires of Ætna could not produce as much in six days as the Cambria iron works turn out in six minutes. Glauber, with all his good sense, taught that the rays of the sun and stars shoot themselves into the earth, and finally became silver and gold. Perhaps he was a prophet, speaking in symbols which he understood not. Now we know that metallurgy does depend on the sun's rays. The sunshine of the carboniferous period has been materialized into coal beds, and now attains perfection in a metal of more real value than gold. In the chemical study of heat, Berthelot's recent work shows culminating progress, and is worthy of him who years ago almost created organic synthesis. After a review of some of the most abstruse speculations in theoretical and physical chemistry, Professor Ordway went on to discuss the importance of biological chemistry. This branch is yet in its infancy, and has few to tenderly care for it. Most chemists prefer to take easier subjects, but the interest in it is increasing. The field is large and there is room for many laborers. Proximate organic analysis still remains undeveloped, and the world does not comprehend the light that we already have. In fermentation, putrefaction, vitrification and zymotic diseases, life may intervene; but how much do we yet know as to what is cause and what is merely concomitant? It is pertinent to ask whether chemistry tends, as many think all physical science tends, to materialism? I believe no true science tends that way; it is the lack of liberal cultivation that leads to such dimness of vision. Materialism is no more prevalent now than among the Athenians, who had no physical science. We hear much of the culture of that people, as if æsthetics were the only science and floriculture the only culture. There is much in the training of the chemist to foster a wholesome skepticism and just intolerance; intolerance of human pride and skepticism of airy theories. In chemical practice the constant appeal to sensible tests and the precision of the balance checks reliance on hasty assumptions. The chemist soon learns that exact truthfulness in others and rigid honesty in himself lie at the very foundation of science and real knowledge; and he looks on laxity in experiment or statement as the unpardonable sin. No other subject is so well calculated to impress one with the idea that theories are but the changeable dress of science. We all wonder what will become of the atomic theory itself when its centennial comes round twenty-seven years hence.

observation at the time. The actual number of species in any one group must always fall far short of the possible number, and for this reason it is out of the question for us to attempt the solution of the problem of derivation, or to hope for any solution beyond one within the most indefinite limits of correctness. If, when we take one of the most limited of the groups of the animal kingdom, we find ourselves engaged in a hopeless task, what must be the prospect should we attack the problem of other classes or groups of the animal kingdom, where the species run into the thousands, while they number only tens in the case we have attempted to follow out? Shall we say "ignorabimus," or "impavidi progrediamur" and valiantly chase a phantom we can never hope to seize?

CHEMISTRY AS AN ART, AND CHEMISTRY AS A SCIENCE.

BY PROFESSOR J. M. ORDWAY.

Professor J. M. Ordway, of Boston, spoke of "Chemistry as an Art, and Chemistry as a Science," comparing both, and pointing out some recent lines of advancement. The past year, he said, has been one of laborious activity in chemistry, but it has not been marked by any epoch-making discoveries. Meyer's recent apparent resolution of the chlorine molecule has not, indeed, been verified by the carefully devised experiments of Crafts, but the latter does seemingly confirm the change of iodine by intense heat. The years 1879 and 1880 will rank hereafter as years in which Meyer found means to throw new light on the nature of the haloids. Twenty-four years ago Perkins sought for artificial quinine, and found instead a better than royal purple. Then, by various hands and in rapid succession, red and yellow and black and brown and blue dyes were brought out from what proved to be something more than aniline. Now the novelty is past, and the announcement of a new dye hardly creates a ripple of excitement. The twelve-year-old synthesis of alizarine has given us colors purer, brighter, faster and cheaper than those of the obsolescent madder. Of late, wool has been provided for, and the extinction of cochineal plantations is threatened by reds of surpassing brilliancy, durability and ease of application. Baeyer has recently effected the synthesis of indigo, and tropical indigo fields may in time share the fate of the madder farms of France and Turkey. But indigo itself will not continue to satisfy our demand. We have become accustomed to hues of a delicacy and richness that no one dared to dream of twenty-five years ago. The æsthetic taste of this generation has been too much pampered; and dyers will soon call for something uniting the brilliancy of the aniline blues with the fixedness of indigo, and its adaptedness to wool and cotton. And Germany which has done the most in studying out these extraordinary colored compounds, now furnished the most of the industrial fruits of seemingly unpractical researches. Investigation costs, investigation pays; in more senses than one our science "opens wide her everdaring gates on golden hinges turning."

The passing years are bringing to light new elementary bodies, and new metals are becoming like new asteroids, of too little mass to influence the orbits of other planets, and too much out of sight to interest many. Within five years fourteen new metals have called for recognition; and in 1879 alone chemists have claimed the discovery of six. Of new alloys, manganian copper is worthy of regard,

since it may in a measure play the part for copper that *spiegeleisen* does for steel.

In 1620 Bacon published the second part of his "Novum Organum," wherein he pointed out the way to appeal to nature by experiment, instead of deriving all science from the teaching of the ancients. But his methods had little immediate influence on the science of the time. He relied on induction; and induction alone simply strings together dry bones. That perception of general principles which makes science comes not altogether from the mere collation of facts. We need something more than eyes to see.

The great chemist of two hundred and fifty years ago was Van Helmont. To him we owe the word gas, which he derived not from *geist*, but from *chaos*, as representing the original form of matter. When our forefathers were laying the foundations of this nation alchemy was in its dotage, and chemistry took its rise in a dim knowledge of the gases. The evolution of chemistry as a science was threefold. First, the study of the gases, then the study of heat, then the study of combining weights. Consider how much of what we now know depends on the gases that Cavendish, Black, Scheele and Priestley revealed. The study of combustion, respiration, vegetable growth, organic decay, geological transformation and hygiene involves the study of carbon dioxide. Carbon monoxide reduces the metals, and plays a part in the Bessemer process for making steel. The fuel of the future is to be coal resolved into a chaos of carbonic oxide and hydrogen. At the end of the last century Murdoch found a use for coal gas, and in its train came a host of secondary products having a marvellous effect on science and industry. A test came into chemistry when Becher attempted to explain combustion. Vulcan of old made as good iron as the blacksmith requires to-day. As for quantity, Vulcan with all his Cyclops and the fires of Ætna could not produce as much in six days as the Cambria iron works turn out in six minutes. Glauber, with all his good sense, taught that the rays of the sun and stars shoot themselves into the earth, and finally became silver and gold. Perhaps he was a prophet, speaking in symbols which he understood not. Now we know that metallurgy does depend on the sun's rays. The sunshine of the carboniferous period has been materialized into coal beds, and now attains perfection in a metal of more real value than gold. In the chemical study of heat, Berthelot's recent work shows culminating progress, and is worthy of him who years ago almost created organic synthesis. After a review of some of the most abstruse speculations in theoretical and physical chemistry, Professor Ordway went on to discuss the importance of biological chemistry. This branch is yet in its infancy, and has few to tenderly care for it. Most chemists prefer to take easier subjects, but the interest in it is increasing. The field is large and there is room for many laborers. Proximate organic analysis still remains undeveloped, and the world does not comprehend the light that we already have. In fermentation, putrefaction, vitrification and zymotic diseases, life may intervene; but how much do we yet know as to what is cause and what is merely concomitant? It is pertinent to ask whether chemistry tends, as many think all physical science tends, to materialism? I believe no true science tends that way; it is the lack of liberal cultivation that leads to such dimness of vision. Materialism is no more prevalent now than among the Athenians, who had no physical science. We hear much of the culture of that people, as if æsthetics were the only science and floriculture the only culture. There is much in the training of the chemist to foster a wholesome skepticism and just intolerance; intolerance of human pride and skepticism of airy theories. In chemical practice the constant appeal to sensible tests and the precision of the balance checks reliance on hasty assumptions. The chemist soon learns that exact truthfulness in others and rigid honesty in himself lie at the very foundation of science and real knowledge; and he looks on laxity in experiment or statement as the unpardonable sin. No other subject is so well calculated to impress one with the idea that theories are but the changeable dress of science. We all wonder what will become of the atomic theory itself when its centennial comes round twenty-seven years hence.

SPECTROSCOPIC NOTES ON OBSERVATIONS— CHIEFLY SOLAR—1879-80.

BY PROF. C. A. YOUNG, of Princeton, N. J.

(a) The magnesium lines of the δ group and the sodium lines have been seen several times (first on June 5, 1880) doubly-reversed in the chromosphere spectrum—*i. e.* a bright line appeared as usual in the centre of the broad dark shade, and then this bright line widened and a thin dark line appeared in its centre. The phenomenon seems to be the exact correlative of the double-reversal of the bright sodium lines observable in the flame of a Bunsen burner under certain circumstances.

(b) I have recently been able to repeat the observations on the H lines first made at Sherman in 1872. In the spectrum of the chromosphere I find both H_1 , and H_2 , (or K, as some call H_2) to be *always* reversed; and what is more, H_1 is *double*, the principal line, which is in the centre of the dark shade, being accompanied by another of about half the strength, one division of Angstrom's scale lower—*i. e.* less refrangible. Since last March I have always been able to observe the two H's whenever I could see h , and H_1 *invariably* double.

In the neighborhood of sun spots however, though both H and K are usually reversed on the solar disc, H_1 is *not* double; its attendant line therefore belongs strictly to the spectrum of the chromosphere, and seems to be identical with No. 271 of my catalogue of chromosphere lines, though its wave length is about 3969 instead of 3970.

The observations were made with grating of 17,280 lines to inch; collimator and telescope 12-inch focus.

(c) A high dispersion spectroscope has been constructed by combining the above-mentioned grating, having nearly four square inches of ruled surface, with collimator and telescope of 3 inches aperture and 42 inches focus, the magnifying power employed varying from 50 to 200. The apparatus is strapped to the tube of the equatorial, and thus kept directed to the sun, an image of which is formed on the slit by an anachromatic lens of 3 inches aperture.

The performance of the grating is admirable when perfectly flat—a force of $\frac{1}{4}$ oz. applied at one corner is however sufficient to distort the plate (of speculum metal) $\frac{3}{8}$ inch thick by about $3\frac{1}{2}$ inches square, to an extent which seriously impairs the definition; it is sensitive to such distortions to a degree entirely unexpected. This instrument doubles an enormous number of the Fraunhofer lines. Out of 47 lines between C and G marked by Thalen as common to the spectra of two or more bodies, 38 are double or triple, 3 are doubtful (from difficulty of identification), and 6 only are single so far as the instrument can show,

(d) Distortion of solar prominences by a diffraction spectroscope. Generally, in such an instrument, the forms seen through the opened slit are either disproportionately extended or compressed along the line of dispersion. If the angle between the normal to the grating and the view-telescope is *less* than that between the normal and the collimator, there will be compression or flattening, and *vice versa*. The mathematical investigation is very simple—

Let n be the order of the spectrum observed.

Let λ be the order of the wave length of the ray.

Let S be the distance between adjacent lines of grating.

Let τ be the angle between normal to grating and telescope.

Let κ be the angle between normal to grating and collimator, and finally $\alpha = \tau + \kappa$ = angle between telescope and collimator, which is supposed constant. Then from the fundamental conditions of spectrum formation $n\lambda = S (\sin. \tau - \sin. \kappa)$ or $\sin. \tau = \frac{n\lambda}{S} + \sin. \kappa$, whence $d\tau = \frac{\cos. \kappa}{\cos. \tau} d\kappa$, or $(\cos. \alpha + \sin. \alpha \tan. \tau) d\kappa$, whence, in general, $d\tau$ will not equal $d\kappa$.

Special cases—

1. If $\kappa = \tau$, there is no distortion—but also no dispersion; it is the case of simple reflection.

2. If $\kappa = 0$, grating being kept normal to the collimator, then $\xi = \sec. \alpha d\kappa$.

3. If $\tau = 0$, grating being kept normal to the telescope and moving with it, then $d\tau = \cos. \alpha d\kappa$.

4. If $\alpha = 90^\circ$ $d\tau = \tan. \tau d\kappa$.

5. If $\alpha = 0$, $d\tau = d\kappa$ and there is no distortion. This is possible only by using the same tube both for collimator and view-telescope, the grating being slightly inclined. The principal difficulty with this form of instrument lies in the reflections from the surface of the object glass, which, it is hoped, may be avoided by a special construction of the lens. An instrument on this plan is in process of construction by the Clarks, for the Physical Laboratory at Princeton, and nearly completed.

ON THE THERMO-ELECTRIC ELECTRO-MOTIVE POWER OF FE. AND PT. IN VACUO.

BY PROF. C. A. YOUNG, of Princeton, N. J.

Eisner, a few months ago, published a paper asserting that the thermo-electric power of Antimony and Bismuth is destroyed by removing them from all contact with oxygen, and inserting them in an atmosphere of pure nitrogen. From this he argues that the thermo-electric force in general is due to the contact of the gases which bathe the metals. The following experiment was tried to test the theory.

By the kindness of Mr. Edison and Mr. Upton a vacuum tube was prepared in Mr. Edison's laboratory, containing an iron wire, about 2 inches long, firmly joined to two platinum terminals which passed through the walls of the tube; the tube was exhausted until a 2-inch induction coil spark would not pass $\frac{1}{8}$ of an inch in the gauge-tube, indicating a residual atmosphere of about one-millionth. The wire was heated too in candescence during the exhaustion, in order to drive off any possible occluded gases. The platinum wires outside the tube were joined to iron wires, the joinings being covered by glass tubes slipped over them, and a sensitive reflecting galvanometer was included in the circuit. By laying the tube and connected joinings in the sunshine, and alternately shading one or several of the joinings, it was found that the electro motive power of the joinings within the tube was precisely the same as that of those without, and the development of current just as rapid. There was no trace of any modification due to the exhaustion.

ON THE ABSOLUTE INVISIBILITY OF ATOMS AND MOLECULES.

BY PROF. A. E. DOLBEAR.

Maxwell gives the diameter of an atom of hydrogen to be such that two millions of them in a row would measure a millimeter, but under ordinary physical conditions most atoms are combined with other atoms to form molecules, and such combinations are of all degrees of complexity; thus a molecule of water contains three atoms, a molecule of alum about one hundred, while a molecule of albumen, according to Mulder, contains nine hundred atoms, and there is no reason to suppose albumen to be the most complex of all molecular compounds. When atoms are thus combined it is fair to assume that they are arranged in the three dimensions of space, and that the diameter of the molecule will be approximately as the cube root of the number of atoms it contains, so that a molecule of alum will be equal to

$$(\sqrt[3]{100} = 4.64) \frac{4.64}{2000000} = \frac{1}{431000} \text{ mm.}$$

and a molecule containing a thousand atoms will have a diameter of $\frac{1}{2000000} = \frac{1}{2000000}$ mm. Now a good microscope, will enable a skilled observer to identify an object so small as the $\frac{1}{40000}$ mm. Beale in his works on the microscope pictures some fungi as minute as that, and Nobert's test bands and the markings upon the *Amphipura pelucida*, which are of about the same degree of fineness, are easily resolved by good lenses. If thus the efficiency of the microscope could be increased fifty times ($\frac{2000000}{40000} = 50$) it

SPECTROSCOPIC NOTES ON OBSERVATIONS— CHIEFLY SOLAR—1879-80.

BY PROF. C. A. YOUNG, of Princeton, N. J.

(a) The magnesium lines of the δ group and the sodium lines have been seen several times (first on June 5, 1880) doubly-reversed in the chromosphere spectrum—*i. e.* a bright line appeared as usual in the centre of the broad dark shade, and then this bright line widened and a thin dark line appeared in its centre. The phenomenon seems to be the exact correlative of the double-reversal of the bright sodium lines observable in the flame of a Bunsen burner under certain circumstances.

(b) I have recently been able to repeat the observations on the H lines first made at Sherman in 1872. In the spectrum of the chromosphere I find both H_1 , and H_2 , (or K, as some call H_2) to be *always* reversed; and what is more, H_1 is *double*, the principal line, which is in the centre of the dark shade, being accompanied by another of about half the strength, one division of Angstrom's scale lower—*i. e.* less refrangible. Since last March I have always been able to observe the two H's whenever I could see h , and H_1 *invariably* double.

In the neighborhood of sun spots however, though both H and K are usually reversed on the solar disc, H_1 is *not* double; its attendant line therefore belongs strictly to the spectrum of the chromosphere, and seems to be identical with No. 271 of my catalogue of chromosphere lines, though its wave length is about 3969 instead of 3970.

The observations were made with grating of 17,280 lines to inch; collimator and telescope 12-inch focus.

(c) A high dispersion spectroscope has been constructed by combining the above-mentioned grating, having nearly four square inches of ruled surface, with collimator and telescope of 3 inches aperture and 42 inches focus, the magnifying power employed varying from 50 to 200. The apparatus is strapped to the tube of the equatorial, and thus kept directed to the sun, an image of which is formed on the slit by an anachromatic lens of 3 inches aperture.

The performance of the grating is admirable when perfectly flat—a force of $\frac{1}{4}$ oz. applied at one corner is however sufficient to distort the plate (of speculum metal) $\frac{3}{8}$ inch thick by about $3\frac{1}{2}$ inches square, to an extent which seriously impairs the definition; it is sensitive to such distortions to a degree entirely unexpected. This instrument doubles an enormous number of the Fraunhofer lines. Out of 47 lines between C and G marked by Thalen as common to the spectra of two or more bodies, 38 are double or triple, 3 are doubtful (from difficulty of identification), and 6 only are single so far as the instrument can show,

(d) Distortion of solar prominences by a diffraction spectroscope. Generally, in such an instrument, the forms seen through the opened slit are either disproportionately extended or compressed along the line of dispersion. If the angle between the normal to the grating and the view-telescope is *less* than that between the normal and the collimator, there will be compression or flattening, and *vice versa*. The mathematical investigation is very simple—

Let n be the order of the spectrum observed.

Let λ be the order of the wave length of the ray.

Let S be the distance between adjacent lines of grating.

Let τ be the angle between normal to grating and telescope.

Let κ be the angle between normal to grating and collimator, and finally $\alpha = \tau + \kappa$ = angle between telescope and collimator, which is supposed constant. Then from the fundamental conditions of spectrum formation $n\lambda = S (\sin. \tau - \sin. \kappa)$ or $\sin. \tau = \frac{n\lambda}{S} + \sin. \kappa$, whence $d\tau = \frac{\cos. \kappa}{\cos. \tau} d\kappa$, or $(\cos. \alpha + \sin. \alpha \tan. \tau) d\kappa$, whence, in general, $d\tau$ will not equal $d\kappa$.

Special cases—

1. If $\kappa = \tau$, there is no distortion—but also no dispersion; it is the case of simple reflection.

2. If $\kappa = 0$, grating being kept normal to the collimator, then $\xi = \sec. \alpha d\kappa$.

3. If $\tau = 0$, grating being kept normal to the telescope and moving with it, then $d\tau = \cos. \alpha d\kappa$.

4. If $\alpha = 90^\circ$ $d\tau = \tan. \tau d\kappa$.

5. If $\alpha = 0$, $d\tau = d\kappa$ and there is no distortion. This is possible only by using the same tube both for collimator and view-telescope, the grating being slightly inclined. The principal difficulty with this form of instrument lies in the reflections from the surface of the object glass, which, it is hoped, may be avoided by a special construction of the lens. An instrument on this plan is in process of construction by the Clarks, for the Physical Laboratory at Princeton, and nearly completed.

ON THE THERMO-ELECTRIC ELECTRO-MOTIVE POWER OF FE. AND PT. IN VACUO.

BY PROF. C. A. YOUNG, of Princeton, N. J.

Eisner, a few months ago, published a paper asserting that the thermo-electric power of Antimony and Bismuth is destroyed by removing them from all contact with oxygen, and inserting them in an atmosphere of pure nitrogen. From this he argues that the thermo-electric force in general is due to the contact of the gases which bathe the metals. The following experiment was tried to test the theory.

By the kindness of Mr. Edison and Mr. Upton a vacuum tube was prepared in Mr. Edison's laboratory, containing an iron wire, about 2 inches long, firmly joined to two platinum terminals which passed through the walls of the tube; the tube was exhausted until a 2-inch induction coil spark would not pass $\frac{1}{8}$ of an inch in the gauge-tube, indicating a residual atmosphere of about one-millionth. The wire was heated too in candescence during the exhaustion, in order to drive off any possible occluded gases. The platinum wires outside the tube were joined to iron wires, the joinings being covered by glass tubes slipped over them, and a sensitive reflecting galvanometer was included in the circuit. By laying the tube and connected joinings in the sunshine, and alternately shading one or several of the joinings, it was found that the electro motive power of the joinings within the tube was precisely the same as that of those without, and the development of current just as rapid. There was no trace of any modification due to the exhaustion.

ON THE ABSOLUTE INVISIBILITY OF ATOMS AND MOLECULES.

BY PROF. A. E. DOLBEAR.

Maxwell gives the diameter of an atom of hydrogen to be such that two millions of them in a row would measure a millimeter, but under ordinary physical conditions most atoms are combined with other atoms to form molecules, and such combinations are of all degrees of complexity; thus a molecule of water contains three atoms, a molecule of alum about one hundred, while a molecule of albumen, according to Mulder, contains nine hundred atoms, and there is no reason to suppose albumen to be the most complex of all molecular compounds. When atoms are thus combined it is fair to assume that they are arranged in the three dimensions of space, and that the diameter of the molecule will be approximately as the cube root of the number of atoms it contains, so that a molecule of alum will be equal to

$$(\sqrt[3]{100} = 4.64) \frac{4.64}{2000000} = \frac{1}{431000} \text{ mm.}$$

and a molecule containing a thousand atoms will have a diameter of $\frac{1}{2000000} = \frac{1}{2000000}$ mm. Now a good microscope, will enable a skilled observer to identify an object so small as the $\frac{1}{40000}$ mm. Beale in his works on the microscope pictures some fungi as minute as that, and Nobert's test bands and the markings upon the *Amphiphura pelucida*, which are of about the same degree of fineness, are easily resolved by good lenses. If thus the efficiency of the microscope could be increased fifty times ($\frac{2000000}{40000} = 50$) it

SPECTROSCOPIC NOTES ON OBSERVATIONS— CHIEFLY SOLAR—1879-80.

BY PROF. C. A. YOUNG, of Princeton, N. J.

(a) The magnesium lines of the δ group and the sodium lines have been seen several times (first on June 5, 1880) doubly-reversed in the chromosphere spectrum—*i. e.* a bright line appeared as usual in the centre of the broad dark shade, and then this bright line widened and a thin dark line appeared in its centre. The phenomenon seems to be the exact correlative of the double-reversal of the bright sodium lines observable in the flame of a Bunsen burner under certain circumstances.

(b) I have recently been able to repeat the observations on the H lines first made at Sherman in 1872. In the spectrum of the chromosphere I find both H_1 , and H_2 , (or K, as some call H_2) to be *always* reversed; and what is more, H_1 is *double*, the principal line, which is in the centre of the dark shade, being accompanied by another of about half the strength, one division of Angstrom's scale lower—*i. e.* less refrangible. Since last March I have always been able to observe the two H's whenever I could see h, and H_1 *invariably* double.

In the neighborhood of sun spots however, though both H and K are usually reversed on the solar disc, H_1 is *not* double; its attendant line therefore belongs strictly to the spectrum of the chromosphere, and seems to be identical with No. 271 of my catalogue of chromosphere lines, though its wave length is about 3969 instead of 3970.

The observations were made with grating of 17,280 lines to inch; collimator and telescope 12-inch focus.

(c) A high dispersion spectroscope has been constructed by combining the above-mentioned grating, having nearly four square inches of ruled surface, with collimator and telescope of 3 inches aperture and 42 inches focus, the magnifying power employed varying from 50 to 200. The apparatus is strapped to the tube of the equatorial, and thus kept directed to the sun, an image of which is formed on the slit by an anachromatic lens of 3 inches aperture.

The performance of the grating is admirable when perfectly flat—a force of $\frac{1}{4}$ oz. applied at one corner is however sufficient to distort the plate (of speculum metal) $\frac{3}{8}$ inch thick by about $3\frac{1}{2}$ inches square, to an extent which seriously impairs the definition; it is sensitive to such distortions to a degree entirely unexpected. This instrument doubles an enormous number of the Fraunhofer lines. Out of 47 lines between C and G marked by Thalen as common to the spectra of two or more bodies, 38 are double or triple, 3 are doubtful (from difficulty of identification), and 6 only are single so far as the instrument can show,

(d) Distortion of solar prominences by a diffraction spectroscope. Generally, in such an instrument, the forms seen through the opened slit are either disproportionately extended or compressed along the line of dispersion. If the angle between the normal to the grating and the view-telescope is *less* than that between the normal and the collimator, there will be compression or flattening, and *vice versa*. The mathematical investigation is very simple—

Let n be the order of the spectrum observed.

Let λ be the order of the wave length of the ray.

Let S be the distance between adjacent lines of grating.

Let τ be the angle between normal to grating and telescope.

Let κ be the angle between normal to grating and collimator, and finally $\alpha = \tau + \kappa$ = angle between telescope and collimator, which is supposed constant. Then from the fundamental conditions of spectrum formation $n\lambda = S (\sin. \tau - \sin. \kappa)$ or $\sin. \tau = \frac{n\lambda}{S} + \sin. \kappa$, whence $d\tau = \frac{\cos. \kappa}{\cos. \tau} d\kappa$, or $(\cos. \alpha + \sin. \alpha \tan. \tau) d\kappa$, whence, in general, $d\tau$ will not equal $d\kappa$.

Special cases—

1. If $\kappa = \tau$, there is no distortion—but also no dispersion; it is the case of simple reflection.

2. If $\kappa = 0$, grating being kept normal to the collimator, then $\xi = \sec. \alpha d\kappa$.

3. If $\tau = 0$, grating being kept normal to the telescope and moving with it, then $d\tau = \cos. \alpha d\kappa$.

4. If $\alpha = 90^\circ$ $d\tau = \tan. \tau d\kappa$.

5. If $\alpha = 0$, $d\tau = d\kappa$ and there is no distortion. This is possible only by using the same tube both for collimator and view-telescope, the grating being slightly inclined. The principal difficulty with this form of instrument lies in the reflections from the surface of the object glass, which, it is hoped, may be avoided by a special construction of the lens. An instrument on this plan is in process of construction by the Clarks, for the Physical Laboratory at Princeton, and nearly completed.

ON THE THERMO-ELECTRIC ELECTRO-MOTIVE POWER OF FE. AND PT. IN VACUO.

BY PROF. C. A. YOUNG, of Princeton, N. J.

Eisner, a few months ago, published a paper asserting that the thermo-electric power of Antimony and Bismuth is destroyed by removing them from all contact with oxygen, and inserting them in an atmosphere of pure nitrogen. From this he argues that the thermo-electric force in general is due to the contact of the gases which bathe the metals. The following experiment was tried to test the theory.

By the kindness of Mr. Edison and Mr. Upton a vacuum tube was prepared in Mr. Edison's laboratory, containing an iron wire, about 2 inches long, firmly joined to two platinum terminals which passed through the walls of the tube; the tube was exhausted until a 2-inch induction coil spark would not pass $\frac{1}{8}$ of an inch in the gauge-tube, indicating a residual atmosphere of about one-millionth. The wire was heated too in candescence during the exhaustion, in order to drive off any possible occluded gases. The platinum wires outside the tube were joined to iron wires, the joinings being covered by glass tubes slipped over them, and a sensitive reflecting galvanometer was included in the circuit. By laying the tube and connected joinings in the sunshine, and alternately shading one or several of the joinings, it was found that the electro motive power of the joinings within the tube was precisely the same as that of those without, and the development of current just as rapid. There was no trace of any modification due to the exhaustion.

ON THE ABSOLUTE INVISIBILITY OF ATOMS AND MOLECULES.

BY PROF. A. E. DOLBEAR.

Maxwell gives the diameter of an atom of hydrogen to be such that two millions of them in a row would measure a millimeter, but under ordinary physical conditions most atoms are combined with other atoms to form molecules, and such combinations are of all degrees of complexity; thus a molecule of water contains three atoms, a molecule of alum about one hundred, while a molecule of albumen, according to Mulder, contains nine hundred atoms, and there is no reason to suppose albumen to be the most complex of all molecular compounds. When atoms are thus combined it is fair to assume that they are arranged in the three dimensions of space, and that the diameter of the molecule will be approximately as the cube root of the number of atoms it contains, so that a molecule of alum will be equal to

$$(\sqrt[3]{100} = 4.64) \frac{4.64}{2000000} = \frac{1}{431000} \text{ mm.}$$

and a molecule containing a thousand atoms will have a diameter of $\frac{1}{2000000} = \frac{1}{2000000}$ mm. Now a good microscope, will enable a skilled observer to identify an object so small as the $\frac{1}{40000}$ mm. Beale in his works on the microscope pictures some fungi as minute as that, and Nobert's test bands and the markings upon the *Amphipura pelucida*, which are of about the same degree of fineness, are easily resolved by good lenses. If thus the efficiency of the microscope could be increased fifty times ($\frac{2000000}{40000} = 50$) it

would be sufficient to enable one to see a molecule of albumen, or if its power could be increased one hundred and seven times it would enable one to see a molecule of alum.

Now Helmholtz has pointed out the probability that interference will limit the visibility of small objects; but suppose that there should be no difficulty from that source, there are two other conditions which will absolutely prevent us from ever seeing the molecule.

1st. Their motions. A free gaseous molecule of hydrogen at the temperature of $0^{\circ}\text{C}.$, and a pressure of 760 mm. mercury, has a free path about $\frac{1}{10000}$ mm. in length, its velocity in this free path being 1860 m. per second or more than a mile, while its direction of movement is changed millions of times per second. Inasmuch as only a glimpse of an object moving no faster than one millimeter per second can be had, for the movements are magnified as well as the object itself, it will be at once seen that a free gaseous molecule can never be seen, not even glimpsed. But suppose such a molecule could be caught and held in the field so it should have no free path. It still has a vibratory motion which constitutes its temperature. The vibratory movement is measured by the number of undulations it sets up in the ether per second, and will average five thousand millions of millions, a motion which would make the space occupied by the molecule visibly transparent, that is it could not be seen. This is true for liquids and solids. Mr. D. N. Hodges finds the path of a molecule of water at its surface to be .0000024 mm., and though it is still much less in a solid it must still be much too great for observation.

2d. They are transparent. The rays of the sun stream through the atmosphere, and the latter is not perceptibly heated by them as it would be if absorption took place in it. The air is heated by conduction, contact with the earth, which has absorbed and transformed the energy of the rays. When selective absorption takes place the number of rays absorbed is small when compared with the whole number presented, so that practically the separate molecules would be too transparent to be seen, though their magnitude and motions were not absolute hindrances.

ON THE AURORA AND ZODIACAL LIGHT OF MAY 2, 1877.

By HENRY C. LEWIS.

A simultaneous appearance of an aurora and the zodiacal light appeared on this evening, and a comparison between them is here given. The various changes of the aurora are given in detail. A remarkable feature was the formation of a bright streamer which maintained its position relative to the earth for nearly an hour. Meanwhile, the Zodiacal Cone, which was bright early in the evening had moved past the streamer and passed below the horizon. The streamer had remained, like the great pointer, fixed to the earth, and marking its motion, while the heavens revolved past it. This fact was conclusive evidence of the terrestrial character of the aurora and of the cosmical character of the zodiacal light. Another fact leading to the same conclusion was the character of their spectra. That of the zodiacal light was continuous, and that of the aurora was a line-spectrum—the former is such as would be given by sunlight reflected from matter in space; the latter would be given by an electric discharge through a gas.

OBSERVATIONS ON BRACHIOPODS.

By PROF. EDW. S. MORSE.

Mr. Morse gave the anatomical details of some Brachiopods he had studied in Japan, and described the existence of a curious parasitic worm in a large species of *Lingula*. He also gave further facts regarding the so-called hearts of certain brachiopods, and expressed his belief that they were glands of some kind connected with the reproductive organs.

THE KAMES OR ESKARS OF MAINE.

By GEO. H. STONE, Kent's Hill, Me.

This paper is accompanied by a map showing the courses of the larger Kame-systems of Maine. Omitting short, isolated ridges of gravel, the map shows thirty distinct systems of Kame gravels, varying from five to one hundred and fifty miles in length. The total length of Kames and Kame-plains thus far mapped is about 2000 miles. The map is the result of amateur explorations made at intervals during the past four years.

The paper discusses the following points regarding the Kames:

1. *Kame drift compared with glacial drift.*

The facts show that Kame material has in general been transported farther than the morainal material which was originally derived from the same locality.

2. *The Kame streams.*

The Kames were deposited by currents flowing lengthwise of their courses, and in all but four undecided cases the currents flowed southwards. The Kame streams resembled sub-aerial rivers in their meanderings, their branches, and in all other respects. All the long systems in the State are much higher at their northern than at their southern ends. The water of these rivers is shown to have flowed faster on long down slopes than on up slopes. There is strong reason to believe that most of the water of the melting glacier escaped by superficial channels, unless near the terminal moraine. Except near the coast there are in Maine almost no signs of sub-glacial streams.

3. *The external forms of Kames.*

1. The single ridge. 2. Reticulated plains, composed of a series of reticulated ridges with enclosed funnels or lakelets. 3. The solid or continuous plains, which are broad, flat-topped ridges, showing few or no signs of separate ridges, and often of great height.

4. *The internal structure of Kames.*

Kames are of two kinds—1. The stratified Kame, which is the more common type. 2. The pell-mell Kame.

The same Kame may be stratified in one part of its course and pell-mell in another.

5. *Action of the sea upon the Kames.*

During the Champlain period the sea stood at a height, in the central parts of Maine, about 300 or 350 feet above the present sea level. The Kames are plainly overlain by the marine clays, and the sea greatly modified their form. The difference between the Kame that has been under the sea and that which has not is often very great, and conclusively proves that the Kames proper cannot have been a marine deposit.

6. *Topographical relations of the Kames.*

No general law of relationship between the Kames and the relief forms of the land can be derived from local observations, for there are many purely local relationships. The only invariable rule thus far established is that the Kames never cross hills more than about 200 feet higher than the country lying to the northward. Maine is traversed by numerous ranges of hills trending eastward or northeastward, and the Kame systems never cross the high ranges except by low passes. Low hills they cross freely. The inference of the writer is that the Kames were deposited when the glacier was so far melted that the higher hills rose above the ice surface, and hence the only escape for the waters southward was by the low passes.

7. *Distribution of the Kames.*

A line joining the northern extremities of the Kames is nearly a straight line; it trends nearly northeast and is roughly parallel with the coast. North and west of this line there are occasional short ridges of Kame origin, but no long systems have yet been discovered.

(The publication of papers read before the recent meetings of the American Association for the Advancement of Science will be continued in our next number.—ED.)

would be sufficient to enable one to see a molecule of albumen, or if its power could be increased one hundred and seven times it would enable one to see a molecule of alum.

Now Helmholtz has pointed out the probability that interference will limit the visibility of small objects; but suppose that there should be no difficulty from that source, there are two other conditions which will absolutely prevent us from ever seeing the molecule.

1st. Their motions. A free gaseous molecule of hydrogen at the temperature of $0^{\circ}\text{C}.$, and a pressure of 760 mm. mercury, has a free path about $\frac{1}{10000}$ mm. in length, its velocity in this free path being 1860 m. per second or more than a mile, while its direction of movement is changed millions of times per second. Inasmuch as only a glimpse of an object moving no faster than one millimeter per second can be had, for the movements are magnified as well as the object itself, it will be at once seen that a free gaseous molecule can never be seen, not even glimpsed. But suppose such a molecule could be caught and held in the field so it should have no free path. It still has a vibratory motion which constitutes its temperature. The vibratory movement is measured by the number of undulations it sets up in the ether per second, and will average five thousand millions of millions, a motion which would make the space occupied by the molecule visibly transparent, that is it could not be seen. This is true for liquids and solids. Mr. D. N. Hodges finds the path of a molecule of water at its surface to be .0000024 mm., and though it is still much less in a solid it must still be much too great for observation.

2d. They are transparent. The rays of the sun stream through the atmosphere, and the latter is not perceptibly heated by them as it would be if absorption took place in it. The air is heated by conduction, contact with the earth, which has absorbed and transformed the energy of the rays. When selective absorption takes place the number of rays absorbed is small when compared with the whole number presented, so that practically the separate molecules would be too transparent to be seen, though their magnitude and motions were not absolute hindrances.

ON THE AURORA AND ZODIACAL LIGHT OF MAY 2, 1877.

By HENRY C. LEWIS.

A simultaneous appearance of an aurora and the zodiacal light appeared on this evening, and a comparison between them is here given. The various changes of the aurora are given in detail. A remarkable feature was the formation of a bright streamer which maintained its position relative to the earth for nearly an hour. Meanwhile, the Zodiacal Cone, which was bright early in the evening had moved past the streamer and passed below the horizon. The streamer had remained, like the great pointer, fixed to the earth, and marking its motion, while the heavens revolved past it. This fact was conclusive evidence of the terrestrial character of the aurora and of the cosmical character of the zodiacal light. Another fact leading to the same conclusion was the character of their spectra. That of the zodiacal light was continuous, and that of the aurora was a line-spectrum—the former is such as would be given by sunlight reflected from matter in space; the latter would be given by an electric discharge through a gas.

OBSERVATIONS ON BRACHIOPODS.

By PROF. EDW. S. MORSE.

Mr. Morse gave the anatomical details of some Brachiopods he had studied in Japan, and described the existence of a curious parasitic worm in a large species of *Lingula*. He also gave further facts regarding the so-called hearts of certain brachiopods, and expressed his belief that they were glands of some kind connected with the reproductive organs.

THE KAMES OR ESKARS OF MAINE.

By GEO. H. STONE, Kent's Hill, Me.

This paper is accompanied by a map showing the courses of the larger Kame-systems of Maine. Omitting short, isolated ridges of gravel, the map shows thirty distinct systems of Kame gravels, varying from five to one hundred and fifty miles in length. The total length of Kames and Kame-plains thus far mapped is about 2000 miles. The map is the result of amateur explorations made at intervals during the past four years.

The paper discusses the following points regarding the Kames:

1. *Kame drift compared with glacial drift.*

The facts show that Kame material has in general been transported farther than the morainal material which was originally derived from the same locality.

2. *The Kame streams.*

The Kames were deposited by currents flowing lengthwise of their courses, and in all but four undecided cases the currents flowed southwards. The Kame streams resembled sub-aerial rivers in their meanderings, their branches, and in all other respects. All the long systems in the State are much higher at their northern than at their southern ends. The water of these rivers is shown to have flowed faster on long down slopes than on up slopes. There is strong reason to believe that most of the water of the melting glacier escaped by superficial channels, unless near the terminal moraine. Except near the coast there are in Maine almost no signs of sub-glacial streams.

3. *The external forms of Kames.*

1. The single ridge. 2. Reticulated plains, composed of a series of reticulated ridges with enclosed funnels or lakelets. 3. The solid or continuous plains, which are broad, flat-topped ridges, showing few or no signs of separate ridges, and often of great height.

4. *The internal structure of Kames.*

Kames are of two kinds—1. The stratified Kame, which is the more common type. 2. The pell-mell Kame.

The same Kame may be stratified in one part of its course and pell-mell in another.

5. *Action of the sea upon the Kames.*

During the Champlain period the sea stood at a height, in the central parts of Maine, about 300 or 350 feet above the present sea level. The Kames are plainly overlain by the marine clays, and the sea greatly modified their form. The difference between the Kame that has been under the sea and that which has not is often very great, and conclusively proves that the Kames proper cannot have been a marine deposit.

6. *Topographical relations of the Kames.*

No general law of relationship between the Kames and the relief forms of the land can be derived from local observations, for there are many purely local relationships. The only invariable rule thus far established is that the Kames never cross hills more than about 200 feet higher than the country lying to the northward. Maine is traversed by numerous ranges of hills trending eastward or northeastward, and the Kame systems never cross the high ranges except by low passes. Low hills they cross freely. The inference of the writer is that the Kames were deposited when the glacier was so far melted that the higher hills rose above the ice surface, and hence the only escape for the waters southward was by the low passes.

7. *Distribution of the Kames.*

A line joining the northern extremities of the Kames is nearly a straight line; it trends nearly northeast and is roughly parallel with the coast. North and west of this line there are occasional short ridges of Kame origin, but no long systems have yet been discovered.

(The publication of papers read before the recent meetings of the American Association for the Advancement of Science will be continued in our next number.—ED.)

would be sufficient to enable one to see a molecule of albumen, or if its power could be increased one hundred and seven times it would enable one to see a molecule of alum.

Now Helmholtz has pointed out the probability that interference will limit the visibility of small objects; but suppose that there should be no difficulty from that source, there are two other conditions which will absolutely prevent us from ever seeing the molecule.

1st. Their motions. A free gaseous molecule of hydrogen at the temperature of $0^{\circ}\text{C}.$, and a pressure of 760 mm. mercury, has a free path about $\frac{1}{10000}$ mm. in length, its velocity in this free path being 1860 m. per second or more than a mile, while its direction of movement is changed millions of times per second. Inasmuch as only a glimpse of an object moving no faster than one millimeter per second can be had, for the movements are magnified as well as the object itself, it will be at once seen that a free gaseous molecule can never be seen, not even glimpsed. But suppose such a molecule could be caught and held in the field so it should have no free path. It still has a vibratory motion which constitutes its temperature. The vibratory movement is measured by the number of undulations it sets up in the ether per second, and will average five thousand millions of millions, a motion which would make the space occupied by the molecule visibly transparent, that is it could not be seen. This is true for liquids and solids. Mr. D. N. Hodges finds the path of a molecule of water at its surface to be .0000024 mm., and though it is still much less in a solid it must still be much too great for observation.

2d. They are transparent. The rays of the sun stream through the atmosphere, and the latter is not perceptibly heated by them as it would be if absorption took place in it. The air is heated by conduction, contact with the earth, which has absorbed and transformed the energy of the rays. When selective absorption takes place the number of rays absorbed is small when compared with the whole number presented, so that practically the separate molecules would be too transparent to be seen, though their magnitude and motions were not absolute hindrances.

ON THE AURORA AND ZODIACAL LIGHT OF MAY 2, 1877.

By HENRY C. LEWIS.

A simultaneous appearance of an aurora and the zodiacal light appeared on this evening, and a comparison between them is here given. The various changes of the aurora are given in detail. A remarkable feature was the formation of a bright streamer which maintained its position relative to the earth for nearly an hour. Meanwhile, the Zodiacal Cone, which was bright early in the evening had moved past the streamer and passed below the horizon. The streamer had remained, like the great pointer, fixed to the earth, and marking its motion, while the heavens revolved past it. This fact was conclusive evidence of the terrestrial character of the aurora and of the cosmical character of the zodiacal light. Another fact leading to the same conclusion was the character of their spectra. That of the zodiacal light was continuous, and that of the aurora was a line-spectrum—the former is such as would be given by sunlight reflected from matter in space; the latter would be given by an electric discharge through a gas.

OBSERVATIONS ON BRACHIOPODS.

By PROF. EDW. S. MORSE.

Mr. Morse gave the anatomical details of some Brachiopods he had studied in Japan, and described the existence of a curious parasitic worm in a large species of *Lingula*. He also gave further facts regarding the so-called hearts of certain brachiopods, and expressed his belief that they were glands of some kind connected with the reproductive organs.

THE KAMES OR ESKARS OF MAINE.

By GEO. H. STONE, Kent's Hill, Me.

This paper is accompanied by a map showing the courses of the larger Kame-systems of Maine. Omitting short, isolated ridges of gravel, the map shows thirty distinct systems of Kame gravels, varying from five to one hundred and fifty miles in length. The total length of Kames and Kame-plains thus far mapped is about 2000 miles. The map is the result of amateur explorations made at intervals during the past four years.

The paper discusses the following points regarding the Kames:

1. *Kame drift compared with glacial drift.*

The facts show that Kame material has in general been transported farther than the morainal material which was originally derived from the same locality.

2. *The Kame streams.*

The Kames were deposited by currents flowing lengthwise of their courses, and in all but four undecided cases the currents flowed southwards. The Kame streams resembled sub-aerial rivers in their meanderings, their branches, and in all other respects. All the long systems in the State are much higher at their northern than at their southern ends. The water of these rivers is shown to have flowed faster on long down slopes than on up slopes. There is strong reason to believe that most of the water of the melting glacier escaped by superficial channels, unless near the terminal moraine. Except near the coast there are in Maine almost no signs of sub-glacial streams.

3. *The external forms of Kames.*

1. The single ridge. 2. Reticulated plains, composed of a series of reticulated ridges with enclosed funnels or lakelets. 3. The solid or continuous plains, which are broad, flat-topped ridges, showing few or no signs of separate ridges, and often of great height.

4. *The internal structure of Kames.*

Kames are of two kinds—1. The stratified Kame, which is the more common type. 2. The pell-mell Kame.

The same Kame may be stratified in one part of its course and pell-mell in another.

5. *Action of the sea upon the Kames.*

During the Champlain period the sea stood at a height, in the central parts of Maine, about 300 or 350 feet above the present sea level. The Kames are plainly overlain by the marine clays, and the sea greatly modified their form. The difference between the Kame that has been under the sea and that which has not is often very great, and conclusively proves that the Kames proper cannot have been a marine deposit.

6. *Topographical relations of the Kames.*

No general law of relationship between the Kames and the relief forms of the land can be derived from local observations, for there are many purely local relationships. The only invariable rule thus far established is that the Kames never cross hills more than about 200 feet higher than the country lying to the northward. Maine is traversed by numerous ranges of hills trending eastward or northeastward, and the Kame systems never cross the high ranges except by low passes. Low hills they cross freely. The inference of the writer is that the Kames were deposited when the glacier was so far melted that the higher hills rose above the ice surface, and hence the only escape for the waters southward was by the low passes.

7. *Distribution of the Kames.*

A line joining the northern extremities of the Kames is nearly a straight line; it trends nearly northeast and is roughly parallel with the coast. North and west of this line there are occasional short ridges of Kame origin, but no long systems have yet been discovered.

(The publication of papers read before the recent meetings of the American Association for the Advancement of Science will be continued in our next number.—ED.)

would be sufficient to enable one to see a molecule of albumen, or if its power could be increased one hundred and seven times it would enable one to see a molecule of alum.

Now Helmholtz has pointed out the probability that interference will limit the visibility of small objects; but suppose that there should be no difficulty from that source, there are two other conditions which will absolutely prevent us from ever seeing the molecule.

1st. Their motions. A free gaseous molecule of hydrogen at the temperature of $0^{\circ}\text{C}.$, and a pressure of 760 mm. mercury, has a free path about $\frac{1}{10000}$ mm. in length, its velocity in this free path being 1860 m. per second or more than a mile, while its direction of movement is changed millions of times per second. Inasmuch as only a glimpse of an object moving no faster than one millimeter per second can be had, for the movements are magnified as well as the object itself, it will be at once seen that a free gaseous molecule can never be seen, not even glimpsed. But suppose such a molecule could be caught and held in the field so it should have no free path. It still has a vibratory motion which constitutes its temperature. The vibratory movement is measured by the number of undulations it sets up in the ether per second, and will average five thousand millions of millions, a motion which would make the space occupied by the molecule visibly transparent, that is it could not be seen. This is true for liquids and solids. Mr. D. N. Hodges finds the path of a molecule of water at its surface to be .0000024 mm., and though it is still much less in a solid it must still be much too great for observation.

2d. They are transparent. The rays of the sun stream through the atmosphere, and the latter is not perceptibly heated by them as it would be if absorption took place in it. The air is heated by conduction, contact with the earth, which has absorbed and transformed the energy of the rays. When selective absorption takes place the number of rays absorbed is small when compared with the whole number presented, so that practically the separate molecules would be too transparent to be seen, though their magnitude and motions were not absolute hindrances.

ON THE AURORA AND ZODIACAL LIGHT OF MAY 2, 1877.

By HENRY C. LEWIS.

A simultaneous appearance of an aurora and the zodiacal light appeared on this evening, and a comparison between them is here given. The various changes of the aurora are given in detail. A remarkable feature was the formation of a bright streamer which maintained its position relative to the earth for nearly an hour. Meanwhile, the Zodiacal Cone, which was bright early in the evening had moved past the streamer and passed below the horizon. The streamer had remained, like the great pointer, fixed to the earth, and marking its motion, while the heavens revolved past it. This fact was conclusive evidence of the terrestrial character of the aurora and of the cosmical character of the zodiacal light. Another fact leading to the same conclusion was the character of their spectra. That of the zodiacal light was continuous, and that of the aurora was a line-spectrum—the former is such as would be given by sunlight reflected from matter in space; the latter would be given by an electric discharge through a gas.

OBSERVATIONS ON BRACHIOPODS.

By PROF. EDW. S. MORSE.

Mr. Morse gave the anatomical details of some Brachiopods he had studied in Japan, and described the existence of a curious parasitic worm in a large species of *Lingula*. He also gave further facts regarding the so-called hearts of certain brachiopods, and expressed his belief that they were glands of some kind connected with the reproductive organs.

THE KAMES OR ESKARS OF MAINE.

By GEO. H. STONE, Kent's Hill, Me.

This paper is accompanied by a map showing the courses of the larger Kame-systems of Maine. Omitting short, isolated ridges of gravel, the map shows thirty distinct systems of Kame gravels, varying from five to one hundred and fifty miles in length. The total length of Kames and Kame-plains thus far mapped is about 2000 miles. The map is the result of amateur explorations made at intervals during the past four years.

The paper discusses the following points regarding the Kames:

1. *Kame drift compared with glacial drift.*

The facts show that Kame material has in general been transported farther than the morainal material which was originally derived from the same locality.

2. *The Kame streams.*

The Kames were deposited by currents flowing lengthwise of their courses, and in all but four undecided cases the currents flowed southwards. The Kame streams resembled sub-aerial rivers in their meanderings, their branches, and in all other respects. All the long systems in the State are much higher at their northern than at their southern ends. The water of these rivers is shown to have flowed faster on long down slopes than on up slopes. There is strong reason to believe that most of the water of the melting glacier escaped by superficial channels, unless near the terminal moraine. Except near the coast there are in Maine almost no signs of sub-glacial streams.

3. *The external forms of Kames.*

1. The single ridge. 2. Reticulated plains, composed of a series of reticulated ridges with enclosed funnels or lakelets. 3. The solid or continuous plains, which are broad, flat-topped ridges, showing few or no signs of separate ridges, and often of great height.

4. *The internal structure of Kames.*

Kames are of two kinds—1. The stratified Kame, which is the more common type. 2. The pell-mell Kame.

The same Kame may be stratified in one part of its course and pell-mell in another.

5. *Action of the sea upon the Kames.*

During the Champlain period the sea stood at a height, in the central parts of Maine, about 300 or 350 feet above the present sea level. The Kames are plainly overlain by the marine clays, and the sea greatly modified their form. The difference between the Kame that has been under the sea and that which has not is often very great, and conclusively proves that the Kames proper cannot have been a marine deposit.

6. *Topographical relations of the Kames.*

No general law of relationship between the Kames and the relief forms of the land can be derived from local observations, for there are many purely local relationships. The only invariable rule thus far established is that the Kames never cross hills more than about 200 feet higher than the country lying to the northward. Maine is traversed by numerous ranges of hills trending eastward or northeastward, and the Kame systems never cross the high ranges except by low passes. Low hills they cross freely. The inference of the writer is that the Kames were deposited when the glacier was so far melted that the higher hills rose above the ice surface, and hence the only escape for the waters southward was by the low passes.

7. *Distribution of the Kames.*

A line joining the northern extremities of the Kames is nearly a straight line; it trends nearly northeast and is roughly parallel with the coast. North and west of this line there are occasional short ridges of Kame origin, but no long systems have yet been discovered.

(The publication of papers read before the recent meetings of the American Association for the Advancement of Science will be continued in our next number.—ED.)

MOUNTAIN SITES FOR ASTRONOMICAL OBSERVATIONS.*

BY PROFESSOR O. STONE.

The question of employing mountain sites for Astronomical Observations is one of considerable importance, and two papers, recently published, add materially to the meagre literature on this subject. The first of these is a report of a series of observations on Mt. Ætna made by Professor Langley during a visit to Europe in the winter of 1878.

The lower portion of Mt. Ætna is described as densely inhabited and covered with rich vegetation. At an elevation of about 2000 feet, however, this suddenly ceases, and above this, with the exception of a few uninhabited plantations of chestnut trees which extend to an altitude of 4000 or 5000 feet, there exist only wastes of lava. At an elevation of some 4200 feet a station was established in a hut built of lava, known as "Casa del Bosco."

Professor Langley remained upon the mountain from Christmas until January 14. His instruments consisted of a $3\frac{1}{4}$ inch telescope loaned by the superintendent of the U. S. Naval Observatory and a spectroscope provided with a Rutherford speculum metal grating of 17,296 lines to the inch, and with collimating and observing telescopes of 1.1 inch aperture and 14 inches focal length.

On clear nights, at ordinary elevations, Professor Langley does not recognize steadily more than six stars in the Pleiades, and on an ordinary clear night at Allegheny he cannot steadily see the companion of Polaris with less than two inches aperture. On Mt. Ætna, however, he could see steadily, notwithstanding the moonlight, nine stars of the Pleiades, with the naked eye, the companion of Polaris, with an aperture of 1.6 inch and 11m. companions of β Leporis and α Tauri, and ϵ and σ Orionis with $3\frac{1}{4}$ inch. From this he concludes that stars of about two thirds the brightness of those visible in England under like telescopic power can be seen on Ætna at the elevation of Casa del Bosco. As far as transparency was concerned, a noticeable advantage over stations at a lower altitude was also shown by observations of the sun.

The results in regard to steadiness of definition were not so satisfactory. Although there was probably less tremor in the stars as seen from Mt. Ætna than would have been the case at a lower station, the difference was not great.

The other paper referred to is the report of Mr. Burnham to the Trustees of the "James Lick Trust" of Observations made on Mt. Hamilton with reference to the location of the Lick Observatory.

Mt. Hamilton seems to have been first suggested as a site for the observatory by Professor Holden in 1874, and afterwards approved by Professor Newcomb. The elevation of the summit is 4250 feet, or only a little greater than that of Professor Langley's station at Casa del Bosco. On this summit Mr. Burnham erected a temporary observatory in which was mounted the magnificent six inch Clark refractor, with which nearly all his double star discoveries had been made. It was equatorially mounted with circles and driving clock. The eye-pieces gave powers up to 400. In addition, a set of meteorological instruments were employed.

Mr. Burnham remained upon Mt. Hamilton just 60 days, from August 17 to October 16, with the exception of three days, September 21-23, spent in San Francisco. As the seeing was first class for the 14 nights immediately preceding the 3 days he was absent, it is fair to presume the same conditions continued. During the

whole time only 11 nights were cloudy or foggy, and of the remaining nights there were 42 when the seeing was *first class* and 7 when it was medium, and *no poor* nights when the sky was clear. Besides obtaining remarkable results in the examination of delicate test objects, a search was made for new doubles, and at the close of the report Mr. Burnham gives a catalogue with observations of 42 such objects, 10 of which have a distance of less than 1". A great many objects were examined by daylight but the air, during the greater part of the day at least, appeared to be no steadier than would be ordinarily found elsewhere.

In conclusion, Mr. Burnham says: "So far as one may judge from the time during which these observations were made, there can be no doubt that Mt. Hamilton offers advantages superior to those found at any point where a permanent observatory has been established. * * * * * The ease with which close pairs can be seen, almost down to the horizon, will be apparent from the southern declination of many of the new double stars. * * * * * Close pairs can be observed at least down to 42° south declination. * * * * *

"What has been said about the advantages of Mt. Hamilton for astronomical purposes, is of course, based upon what was seen during the time spent on the mountain. This was my first visit to the Pacific coast, and hence I have no personal knowledge concerning other seasons of the year. From inquiries in various quarters I am satisfied there was nothing about this season unusual, and there seems to be every reason for supposing, as the same cloudless sky and dry air prevails from about March until the commencement of the rainy season, near the close of the year, that the whole of this interval would be equally favorable for the use of the telescope."

One of the most remarkable and interesting conditions observed was the dryness of the atmosphere. The average difference between the wet and dry bulb thermometers was 18°.4 during the first five weeks of Mr. Burnham's residence on Mt. Hamilton and every night was first class when this difference reached 15° and upwards.

Notwithstanding, however, the advantages of a mountain site for an observatory, there are many drawbacks. Even the loneliness of the situation is a disadvantage to the greatest activity. Taking everything into consideration, therefore, probably as favorable a location as any for the next great American observatory is to be found on the plateaus of Colorado.

We have received the following publications from the U. S. Department of the Interior (Bureau of Education).

THE INDIAN SCHOOL AT CARLISLE BARRACKS which acquaints educators and school officials with the interesting experiments of training Indian children in the knowledge and usages of civilized life, in progress during the past eight months at Carlisle Barracks.

VACATION COLONIES FOR SICKLY SCHOOL CHILDREN This subject has for some time received the attention of this department, and as early as 1872 papers by that distinguished and benevolent physician, J. M. Toner, M. D., of Washington, were published, advocating free camping grounds or parks, where poor children and their parents could lodge during the excessive heat of summer.

PROGRESS OF WESTERN EDUCATION IN CHINA AND SIAM. This is an interesting account of the progress of Western ideas and educational methods in China and Siam, forwarded to the department by the United States Minister at Peking, and the United States Consul at Bangkok.

LEGAL RIGHTS OF CHILDREN. This is an elaborate report covering nearly a hundred pages, and treats of the rights of children in the various States of the Union, including education, and also a comparative view of the systems of education in the different States established to give force and effect to those rights, and thus assure the welfare of the individual and the State.

*Observations on Mount Etna, by S. P. Langley. From the *American Journal of Science*, Vol. XIX, July, 1880.

Report to the Trustees of the "James Lick Trust," of Observations made on Mt. Hamilton with reference to the location of Lick Observatory; by S. W. Burnham. Chicago, 1880.